

Self-adjoint codimension 2 boundary conditions for Dirac operators

Bernd Ammann¹ Nadine Große²

¹Universität Regensburg, Germany

²Universität Freiburg, Germany

Section: **Differential Geometry and Global Analysis**

Organized by: Vicente Cortés and Oliver Goertsches

Annual Meeting of the DMV in Chemnitz (virtual)

Sept 15, 2020

Slides available on

<http://www.mathematik.uni-regensburg.de/ammann/talks/2020DMV-handout.pdf>



Table of Contents

Classical boundary conditions for the Laplace operator

- Some classical problems

- The minimal and the maximal operator

- Symplectic structure on boundary data

- Self-adjoint extensions

Codimension 1 bdy cond. for Dirac operators

- How to adapt this for Dirac operators?

- Self-adjoint extensions for D

Codimension 2 bdy cond. for Dirac operators

- The setting of our work

- Monodromy

- Main examples

- Results

- Portman-Sok-Solovej-boundary condition

- Example: 2-dimensional model space

Classical boundary conditions for the Laplace operator

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary.
To model problems in physics or technical applications we often want to solve the **Poisson problem**

$$\Delta u = f$$

for a given function $f \in C^\infty(\Omega)$ or the **eigenvalue problem**

$$\Delta u = \lambda u$$

for $\lambda \in \mathbb{R}$.

Many solutions, and many solutions behave unphysically at the boundary.

↪ We need boundary conditions!

The minimal and the maximal operator

We define

$$\Delta_c : C_c^\infty(\Omega) \rightarrow C_c^\infty(\Omega)$$

as a densely defined operator in $L^2(\Omega)$.

The **minimal operator** Δ_{\min} is the closure of Δ_c with respect to the graph norm.

$$L^2(\Omega) \supset H_0^2(\Omega) \xrightarrow{\Delta_{\min}} L^2(\Omega)$$

And the **maximal operator** is defined as its adjoint

$$\underbrace{\left\{ u \in L^2(\Omega) \mid \Delta u \in L^2(\Omega) \right\}}_{:= \text{dom}(\Delta_{\max}) \supseteq H^2(\Omega)} \xrightarrow{\Delta_{\max}} L^2(\Omega).$$

The map

$$\begin{aligned} \tau = (\tau_D, \tau_N) : C^\infty(\overline{\Omega}) &\rightarrow C^\infty(\partial\Omega) \times C^\infty(\partial\Omega), \\ u &\mapsto (u|_{\partial\Omega}, (\partial_\nu u)|_{\partial\Omega}) \end{aligned}$$

extends to a **trace map** $\tau : \text{dom}(\Delta_{\max}) \rightarrow \check{H}(\partial\Omega)$.



Symplectic structure on boundary data

Trace map $\tau : \text{dom}(\Delta_{\max}) \rightarrow \check{H}(\partial\Omega)$.

Here $\check{H}(\partial\Omega)$ is a Sobolev type Hilbert space,

$$H^{3/2}(\partial\Omega) \times H^{1/2}(\partial\Omega) \subset \check{H}(\partial\Omega) \subset H^{-1/2}(\partial\Omega) \times H^{-3/2}(\partial\Omega)$$

on which a non-degenerate perfect symplectic pairing

$$\begin{aligned} \check{H}(\partial\Omega) \times \check{H}(\partial\Omega) &\xrightarrow{B} \mathbb{R} \\ \left((v_1, w_1), (v_2, w_2) \right) &\mapsto \int_{\partial\Omega} (v_1 w_2 - v_2 w_1) \end{aligned}$$

is well-defined.

Green identity

$$\int_{\Omega} (\Delta u_1) u_2 - \int_{\Omega} u_1 \Delta u_2 = \pm B(\tau(u_1), \tau(u_2)).$$

Self-adjoint extensions

If $A \subset \check{H}(\partial\Omega)$ is a closed subspace, then

$$\underbrace{\tau^{-1}(A)}_{\subset \text{dom}(\Delta_{\max})} \xrightarrow{\Delta_A := \Delta_{\max}|_{\tau^{-1}(A)}} L^2(\Omega)$$

is a closed extension.

Δ_A is self-adjoint iff A is a Lagrangian subspace of $(\check{H}(\partial\Omega), B)$.

Examples:

Dirichlet boundary conditions: $A := \{(0, w) \in \check{H}(\partial\Omega)\}$

Neumann boundary conditions: $A := \{(v, 0) \in \check{H}(\partial\Omega)\}$

Under some regularity assumptions:

- ▶ solution of the Poisson problem
- ▶ discrete, real spectrum
- ▶ eigenspaces finite-dimensional, spanned by smooth functions

Codimension 1 bdy cond. for Dirac operators

Now replace:

$\bar{\Omega} \subset \mathbb{R}^n$	$C^\infty(\bar{\Omega})$	Laplacian Δ
cpct. Riem. spin manfd. with boundary M	twisted spinors $C^\infty(M, V \otimes \Sigma M)$	twisted Dirac oper. \not{D}

$$\begin{aligned} \tau =: C^\infty(M, V \otimes \Sigma M) &\rightarrow C^\infty(\partial M, (V \otimes \Sigma M)|_{\partial M}), \\ u &\mapsto u|_{\partial \Omega} \end{aligned}$$

extends to a **trace map** $\tau : \text{dom}(\not{D}_{\max}) \rightarrow \check{H}(\partial M)$.

Here $\check{H}(\partial \Omega)$ is a Sobolev type Hilbert space,

$$H^{1/2}(\partial M, (V \otimes \Sigma M)|_{\partial M}) \subset \check{H}(\partial M) \subset H^{-1/2}(\partial M, (V \otimes \Sigma M)|_{\partial M})$$

with a skew-hermitian sesquilinear map

$$B : \check{H}(\partial M) \times \check{H}(\partial M) \rightarrow \mathbb{C}.$$

$$\int_M \langle \not{D}\varphi, \psi \rangle - \int_M \langle \varphi, \not{D}\psi \rangle = B(\tau(\varphi), \tau(\psi)).$$

Self-adjoint extensions for D

If $A \subset \check{H}(\partial M)$ is a closed subspace, then

$$\underbrace{\tau^{-1}(A)}_{\subset \text{dom}(\not{D}_{\max})} \xrightarrow{\not{D}_A := \not{D}_{\max}|_{\tau^{-1}(A)}} L^2(M, V \otimes \Sigma M)$$

is a closed extension.

\not{D}_A is self-adjoint iff A is a Lagrangian subspace of $(\check{H}(\partial M), B)$.

Under some regularity assumptions:

- ▶ discrete, real spectrum
- ▶ eigenspaces finite-dimensional, spanned by smooth sections
- ▶ \not{D}_A is a Fredholm operator

\rightsquigarrow Atiyah-Patodi-Singer index theorem.

See e.g. Bär-Ballmann [arxiv 1101.1196](https://arxiv.org/abs/1101.1196).

The setting of our work

- ▶ Let (M, g) be a complete oriented Riemannian manifold, N a compact oriented submanifold of codimension 2.
- ▶ We assume that $M \setminus N$ is spin. Thus there is a complex spinor bundle $\Sigma \rightarrow M \setminus N$.
- ▶ Let $L \rightarrow M \setminus N$ be a flat hermitian line bundle. We get

$$\pi_1(M \setminus N) \rightarrow S^1 \subset \mathbb{C}$$

- ▶ $W := \Sigma \otimes L$ generalized spinor bundle over $M \setminus N$

More general frameworks are possible which will not be discussed in this talk.


Monodromy

Monodromy $\alpha = (\alpha_1, \dots, \alpha_j)$.

$N = \prod_{j=1}^{\ell} N_j$ decomposition into connected components

Parallel transport in W around N_j is $e^{2\pi i \alpha_j}$.

$[\alpha_j] \in \mathbb{R}/\mathbb{Z}$ only depends on j .



Main examples

- ▶ M spin. Monodromy comes from L .
Main subcase: L flat. Monodromy $\pi_1(M \setminus N) \rightarrow S^1$.
Main subsubcase: N is a link in S^3 .

$$(S^1)^\ell \ni (\exp 2\pi i \alpha_j)_{j \in \{1, \dots, \ell\}} \mapsto L_\alpha$$

- ▶ M is not spin, (more precisely: spin structure does not extend), N connected.
Then monodromy only comes from Σ , $\alpha = 1/2 \pmod{\mathbb{Z}}$.
Main subcase: $L = \mathbb{C}$
Example: $M = \mathbb{C}P^{2r}$, $N = \mathbb{C}P^{2r-1}$.
Fix $p \in M \setminus N$, solve $\not{D}\Psi = \psi_0 \delta_p$ on $M \setminus N$ with bdy cond.
Expectation: If PMT would fail, we would get a map

$$S(\Sigma_p) \times \{\text{bdy cond}\} \rightarrow \{\text{non-zero spinors on } N\}.$$

Interesting applications?



Results by Ammann and Große

We obtain

- ▶ a bundle $S \rightarrow N$,
- ▶ a Hilbert space of sections $\check{H}(N, S)$,
- ▶ a skew-hermitian sesquilinear non-degenerate perfect pairing $B : \check{H}(N, S) \times \check{H}(N, S) \rightarrow \mathbb{C}$
- ▶ a trace map $\tau : \text{dom}(\not{D}_{\max}) \rightarrow \check{H}(N, S)$

If $A \subset \check{H}(N, S)$ is a closed subspace, then

$$\underbrace{\tau^{-1}(A)}_{\subset \text{dom}(\not{D}_{\max})} \xrightarrow{\not{D}_A := \not{D}_{\max}|_{\tau^{-1}(A)}} L^2(M, V \otimes \Sigma M)$$

is a closed extension.

$\text{dom}(\not{D}_{\emptyset}) = \text{dom}(\not{D}_{\min})$ and $\text{dom}(\not{D}_{\check{H}(N, S)}) = \text{dom}(\not{D}_{\max})$.

\not{D}_A is self-adjoint iff A is a Lagrangian subspace of $(\check{H}(\partial M), B)$.

Some remarks

- ▶ In the case $\alpha \in \mathbb{Z}$ the submanifold N is invisible, i.e.

$$\mathcal{D}_{\min} = \mathcal{D}_{\max} = \mathcal{D}^M$$

- ▶ τ is not the extension of a restriction map.
In contrast we have: Suppose that $\varphi \in \text{dom}(\mathcal{D}_{\max})$ is bounded on a neighbourhood of N . Then $\varphi \in \text{dom}(\mathcal{D}_{\min})$.
- ▶ The bundle $S \rightarrow N$ has a Clifford multiplication

$$TM|_N \otimes S \rightarrow S$$

Normal volume element: $\omega_{\text{nor}} := e_1 \cdot e_2$ if (e_1, e_2) is a positively oriented orthonormal basis of the normal bundle. This gives a splitting

$$S = S^+ \oplus S^-$$

into the $\pm i$ -eigenspace bundles for the Clifford action of ω_{nor} .



Example: Portman & Sok & Solovej

$\check{H}(N, S^+)$ and $\check{H}(N, S^-)$ are Lagrangian subspaces of $\check{H}(N, S)$.

PSS (2015–2018) considered the special case that $M = S^3$ is the round sphere and N is a link.

Electrons coupled magnetic fields.

Existence of harmonic spin^c-spinors yield statements of the type

If our world is stable, then the fine structure constant $\hbar c/e^2$ has to satisfy some bounds.

Measurements: $\hbar c/e^2 = 137.03599968 \dots$

A spectral flow argument yields harmonic spinors.

Question: How is this spectral flow related to classical link invariants?

2-dimensional model space

Assume $M = \mathbb{C} \ni z$, $N = \{0\}$, $\Sigma = \underline{\mathbb{C}^2} = \Sigma_+ \oplus \Sigma_-$

L flat bundle over $[\mathbb{C} : \{0\}]$, monodromy α

Then ω_{nor} is the standard volume element.

$$\not{D} = \not{D}^{\text{nor}} = \sqrt{2} \begin{pmatrix} 0 & \bar{\partial} \\ -\partial & 0 \end{pmatrix}$$

$\frac{z^{-\alpha}}{|z|^{-\alpha}}$ represents a nowhere vanishing smooth section of L .

Ansatz:

$$\Phi_{\beta,\gamma}^+ := \begin{pmatrix} z^\beta \bar{z}^\gamma \\ 0 \end{pmatrix}, \quad \Phi_{\beta,\gamma}^- := \begin{pmatrix} 0 \\ z^\beta \bar{z}^\gamma \end{pmatrix}.$$

where β and γ over real numbers with $\beta - \gamma + \alpha \in \mathbb{Z}$.

$\Phi_{\beta,\gamma}^\pm \in L_{\text{loc}}^2$ iff $\beta + \gamma > -1$

$$\not{D}\Phi_{\beta,\gamma}^+ = -\sqrt{2}\beta\Phi_{\beta-1,\gamma}^-, \quad \not{D}\Phi_{\beta,\gamma}^- = \sqrt{2}\gamma\Phi_{\beta,\gamma-1}^+$$

Lemma.

The condition that $\Phi_{\beta,\gamma}^{\pm} \in \text{dom}(\mathcal{D}_{\max})$ is characterized as follows (“locally around 0”).

- (1) Suppose $\beta \neq 0$ and $\gamma \neq 0$. Then $\Phi_{\beta,\gamma}^{\pm} \in \text{dom}(\mathcal{D}_{\max})$ if and only if $\beta + \gamma > 0$.
- (2) Suppose $\beta = 0$ and $\gamma \neq 0$. Then $\Phi_{0,\gamma}^{+} \in \text{dom}(\mathcal{D}_{\max})$ if and only if $\gamma > -1$, and $\Phi_{0,\gamma}^{-} \in \text{dom}(\mathcal{D}_{\max})$ if and only if $\gamma > 0$.
- (3) Suppose $\beta \neq 0$ and $\gamma = 0$. Then $\Phi_{\beta,0}^{+} \in \text{dom}(\mathcal{D}_{\max})$ if and only if $\beta > 0$, and $\Phi_{\beta,0}^{-} \in \text{dom}(\mathcal{D}_{\max})$ if and only if $\beta > -1$.
- (4) Suppose $\beta = \gamma = 0$. $\Phi_{0,0}^{\pm} \in \text{dom}(\mathcal{D}_{\max}) = \text{dom}(\mathcal{D}_{\min})$.

$\alpha \in (0, 1)$: Then elements in $\text{dom}(\mathcal{D}_{\max})$ are of the form

$$\left(\begin{array}{c} \bar{z}^{\alpha-1} \varphi_{+} \\ z^{-\alpha} \varphi_{-} \end{array} \right) + \text{dom}(\mathcal{D}_{\min}).$$

Disclaimer

The project is still work in progress. Until now we have only written up in detail the case of a totally geodesic submanifold N and some similar assumptions, but we do not expect any modifications for the general case. Preprints are not yet available.

Summary

- ▶ We have a complete description of the self-adjoint extensions of twisted Dirac operators on manifolds $M^m \setminus N^{m-2}$.
- ▶ The space N is invisible, if the monodromy is trivial.
- ▶ Spinors in the domain of the maximal operator which are not in the minimal domain are not bounded (near the boundary). The trace map has to be modified considerably.
- ▶ Boundary data are given by elements in a Hilbert space \check{H} of sections of a bundle $S \rightarrow N$. This space depends strongly on the monodromy.

Thanks...

... to Boris Botvinnik and Nikolai Saveliev for discussions about link invariants associated to these results

