

$SU(2)$ -CYCLIC SURGERIES AND THE PILLOWCASE

STEVEN SIVEK AND RAPHAEL ZENTNER

ABSTRACT. We study knots in S^3 with infinitely many $SU(2)$ -cyclic surgeries, which are Dehn surgeries such that every representation of the resulting fundamental group into $SU(2)$ has cyclic image. We show that for every such nontrivial knot K , its set of $SU(2)$ -cyclic slopes is bounded and has a unique limit point, which is both a rational number and a boundary slope for K . We also show that such knots are prime and have infinitely many instanton L-space surgeries. Our methods include the application of holonomy perturbation techniques to instanton knot homology, using a strengthening of recent work by the second author.

1. INTRODUCTION

The cyclic surgery theorem of Culler, Gordon, Luecke, and Shalen [CGLS87] says that if a knot in S^3 is neither the unknot nor a torus knot, and if both r - and s -surgery on that knot give manifolds with cyclic fundamental group, then the distance $\Delta(r, s)$ between these slopes is at most 1, and hence there are at most two such nontrivial surgeries. This is the strongest possible result: Moser [Mos71] showed that torus knots have infinitely many lens space surgeries, and for example the pretzel knot $P(-2, 3, 7)$ has two lens space surgeries, of slopes 18 and 19.

In this paper we study a weaker property of surgeries on knots in S^3 . Following Lin [Lin16], we say that a 3-manifold Y is $SU(2)$ -cyclic if all representations $\pi_1(Y) \rightarrow SU(2)$ have cyclic image. Kronheimer and Mrowka [KM04a] proved that any $SU(2)$ -cyclic surgery of slope r on a nontrivial knot $K \subset S^3$ satisfies $|r| > 2$, and Lin [Lin16] proved that any two $SU(2)$ -cyclic slopes $r_i = \frac{p_i}{q_i}$ ($i = 1, 2$) for K satisfy the inequality $\Delta(r_1, r_2) \leq |p_1| + |p_2|$. We will call a nontrivial knot $SU(2)$ -averse if it has infinitely many $SU(2)$ -cyclic surgeries.

If K is $SU(2)$ -averse, then the slopes of its $SU(2)$ -cyclic surgeries form an infinite subset of $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$, so they must have a limit point. One can deduce from Lin's inequality or from Theorem 3.2 below that this limit point is unique; we call this point the *limit slope* of K . For example, the torus knot $T_{p,q}$ has limit slope pq , since $(pq + \frac{1}{n})$ -surgery on $T_{p,q}$ is a lens space for all integers $n \neq 0$.

By studying the $SU(2)$ character variety of the exterior $S^3 \setminus N(K)$, and its image under restriction in the character variety of the peripheral torus $\partial N(K)$, we prove the following.

Theorem 1.1. *Let $K \subset S^3$ be an $SU(2)$ -averse knot. Then its limit slope $r(K)$ is a rational number, with $|r(K)| > 2$, and it is a boundary slope for K . Moreover, K is prime, and the manifold $S^3_s(K)$ constructed by s -surgery on K is an instanton L-space for all rational $s \geq \lceil r(K) \rceil - 1$ if $r(K) > 0$, or for all $s \leq \lfloor r(K) \rfloor + 1$ if $r(K) < 0$.*

Corollary 1.2. *Let K be a nontrivial knot in S^3 . Then there is a constant N such that there is an irreducible representation*

$$\pi_1(S^3_r(K)) \rightarrow SU(2)$$

for all rational r with $|r| > N$.

Proof. If the set of $SU(2)$ -cyclic slopes for a knot K is unbounded, then K must be $SU(2)$ -averse with limit slope ∞ , contradicting $r(K) \in \mathbb{Q}$. \square

In Theorem 1.1, the finiteness and rationality of $r(K)$ are proved as Theorems 4.2 and 5.4 respectively, and the bound $|r(K)| > 2$ comes from Theorem 3.2 (though $r(K) \geq 2$ is already implied by [KM04a]). The fact that $r(K)$ is a boundary slope is Theorem 6.4; this means that there is a properly embedded, essential surface in $S^3 \setminus N(K)$ whose boundary is a nonempty union of curves of slope $r(K)$. The primeness of K is proved as Theorem 10.4. The assertion about instanton L-spaces, namely that for each $s = \frac{a}{b}$ in the given range the singular instanton knot homology $I^\#(S^3_{a/b}(K), \emptyset)$ defined in [KM11] has rank $|a|$, is Theorem 9.1.

We can also prove something about the set of $SU(2)$ -cyclic surgery slopes of an $SU(2)$ -averse knot. Combining Theorem 3.2 and Theorem 8.2, we have:

Theorem 1.3. *Let K be an $SU(2)$ -averse knot with limit slope $r(K)$. Then all $SU(2)$ -cyclic slopes $\frac{m}{n}$ satisfy*

$$\left| \frac{m}{n} - r(K) \right| \leq \frac{|r(K)|}{n},$$

and there is an integer $N \geq 1$ such that $r(K) + \frac{1}{kN}$ is an $SU(2)$ -cyclic slope for all but finitely many $k \in \mathbb{Z}$.

A knot $K \subset S^3$ is said to be *small* if there are no closed, incompressible surfaces in its exterior other than boundary-parallel tori; examples include all two-bridge knots [HT85] and Montesinos knots with at most three rational tangles [Oer84]. In this case we know even more, as proved in Theorems 7.1 and 7.3 and Proposition 7.5.

Theorem 1.4. *If K is a small, $SU(2)$ -averse knot with Alexander polynomial $\Delta_K(t) = \sum_{j=-d}^d a_j t^j$, then the limit slope $r(K)$ is an integer satisfying $|r(K)| \geq 6$ and*

$$|r(K)| \geq 1 + \sum_{j=-d}^d |a_j|.$$

In this case, all but finitely many conjugacy classes of irreducible representations

$$\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$$

satisfy $\rho(\mu^{r(K)} \lambda) = \pm I$, where μ and λ are a meridian and longitude of K .

Finally, by studying how $SU(2)$ -averseness behaves under satellite operations we can also say something about the potential existence of $SU(2)$ -averse knots other than torus knots. Notably, we show in Proposition 10.5 that if $r(K) = \frac{p}{q}$ with $q \geq 2$, then the (p, q) -cable of K is $SU(2)$ -averse with limit slope pq . However, we conjecture that this never happens:

Conjecture 1.5. *If K is a nontrivial $SU(2)$ -averse knot, then K is a torus knot.*

Indeed, in Section 11, we verify Conjecture 1.5 for all alternating Montesinos knots with at most three rational tangles (including all two-bridge knots) and for all knot types through 10 crossings except possibly 10₉₈. The conjecture also holds for algebraic knots.

Outline. The proof of Theorem 1.1 comes from studying the image of the $SU(2)$ character variety of $\pi_1(S^3 \setminus N(K))$ in the *pillowcase*, which is the $SU(2)$ character variety of $\pi_1(\partial N(K))$. Up to conjugacy, every representation $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ satisfies

$$\rho(\mu) = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{-i\alpha} \end{pmatrix}, \quad \rho(\lambda) = \begin{pmatrix} e^{i\beta} & 0 \\ 0 & e^{-i\beta} \end{pmatrix}$$

for some constants α and β , and hence determines a unique point (α, β) of the pillowcase

$$P \cong \frac{(\mathbb{R}/2\pi\mathbb{Z}) \times (\mathbb{R}/2\pi\mathbb{Z})}{(\alpha, \beta) \sim (-\alpha, -\beta)}.$$

A slope $\frac{m}{n}$ is $SU(2)$ -cyclic precisely when there are no images of irreducible characters of $\pi_1(S^3 \setminus K)$ along the line $m\alpha + n\beta = 0$ in P .

Our key observation, developed in Section 3, is that the line $m\alpha + n\beta = 0$ can only avoid a given non-constant path $\gamma : [0, 1] \rightarrow P$ through images of irreducibles if $-\frac{m}{n}$ is a reasonable Diophantine approximation to the slope of the line from $\gamma(0)$ to $\gamma(1)$. We deduce that if K is $SU(2)$ -averse, then any such path must be a straight line of slope $-r(K)$. (Conversely, we show in Section 8 that if every such path is a straight line of rational slope $-r$, then K is $SU(2)$ -averse with $r(K) = r$.)

In Section 4, we show that $r(K) \neq \infty$ by finding (in nearly all cases) an arc of images of irreducibles which is not a vertical line. We do this by making use of Kronheimer and Mrowka’s instanton knot homology [KM10b], or KHI , which by construction is closely related to the space of irreducible representations $\pi_1(S^3 \setminus K) \rightarrow SU(2)$ which send a meridian to a traceless matrix. Using a strengthened version of recent work of the second author [Zen16], we can perturb the Chern-Simons functional which defines $KHI(K)$ so that if K has $SU(2)$ -cyclic surgeries and no such arc of irreducibles, then $KHI(K)$ is the same as KHI of the unknot, which means that K must have been unknotted. More precisely, we prove the following.

Theorem 4.1. *Let K be a nontrivial knot in S^3 . Then at least one of the following is true:*

(1) *For every $\beta \in \mathbb{R}/2\pi\mathbb{Z}$, there is an irreducible $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ such that*

$$\rho(\mu) = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad \text{and} \quad \rho(\lambda) = \begin{pmatrix} e^{i\beta} & 0 \\ 0 & e^{-i\beta} \end{pmatrix}.$$

(2) *There is a constant $\varepsilon > 0$ and an arc of irreducible representations*

$$\rho_s : \pi_1(S^3 \setminus K) \rightarrow SU(2), \quad s \in \left[\frac{\pi}{2}, \frac{\pi}{2} + \varepsilon \right)$$

such that $\rho_s(\mu) = \text{diag}(e^{is}, e^{-is})$ for all s .

This may be viewed as a strengthening of [KM10b, Corollary 7.17]; see also related work of Herald [Her97], Collin–Steer [CS99], or Heusener–Kroll [HK98], in which similar results are proved under assumptions on the equivariant signatures of K .

Remark 1.6. If K is small then the second case of Theorem 4.1 always applies, since the other case is ruled out by first arguing as in the proof of Theorem 6.3 that it would force ∞ to be a boundary slope and then applying [CGLS87, Theorem 2.0.3].

We mention here the needed strengthening of the approximation result in [Zen16], which is of independent interest. Recall that a shearing isotopy (ζ_t) on the 2-dimensional torus is an isotopy through area-preserving maps of the form

$$\zeta_t : (x, y) \mapsto (x, y) + tf(w \cdot (x, y))v, \tag{1.1}$$

where $v, w \in \mathbb{Z}^2$ are orthogonal vectors with w nonzero, $f : \mathbb{R} \rightarrow \mathbb{R}$ is a 2π -periodic function, and \cdot denotes the standard inner product on \mathbb{R}^2 . We say that an isotopy (ϕ_t) is a piecewise isotopy through shearings if there is a partition $0 = t_0 < t_1 < \dots < t_n = 1$ such that for $t_i \leq t \leq t_{i+1}$, the map ϕ_t is given as

$$\phi_t = \zeta_{t-t_i}^{(i)} \circ \phi_{t_i}$$

for some shearing isotopy $(\zeta_s^{(i)})_{s \in [0,1]}$. For fixed $p \in T^2$, the path $t \mapsto \phi_t(p)$ of a piecewise isotopy through shearings is continuous but need not be smooth.

Theorem 4.5. *Let $r \geq 0$. Let $(\psi_t)_{t \in [0,1]}$ be an isotopy through area-preserving maps $\psi_t : T^2 \rightarrow T^2$, and let $\varepsilon > 0$. Then there is a piecewise isotopy through shearings $(\phi_t)_{t \in [0,1]}$ such that ϕ_t is ε -close to ψ_t in the C^r -topology for all $t \in [0, 1]$.*

The rationality of $r(K)$ follows in Section 5 from some foundational results in real algebraic geometry. Using the fact that the pillowcase image of the $SU(2)$ character variety is a semi-algebraic set defined over the field of real algebraic numbers, and hence many of its points have algebraic coordinates, we manage to prove that a straight line in this image must have rational slope. This allows us to further show that if $r(K) = \frac{p}{q}$, then along any such path of representations ρ , there is a real constant c such that

$$\rho(\mu^p \lambda^q) = \begin{pmatrix} e^{ic} & 0 \\ 0 & e^{-ic} \end{pmatrix}$$

up to conjugacy, where e^{ic} is an algebraic number.

In Section 6, we study $SU(2)$ -averse knots via the A -polynomial, which describes the image of the $SL_2(\mathbb{C})$ character variety of $\pi_1(S^3 \setminus K)$ in the $SL_2(\mathbb{C})$ character variety of the peripheral torus. The results of Section 3 imply that if $r(K) = \frac{p}{q}$ then $A_K(M, L)$ is a multiple of $M^p L^q - e^{ic}$, and hence by applying deep results from [CCG⁺94] about the Newton polygon and edge polynomials of $A_K(M, L)$, we see that $r(K)$ is a boundary slope and that e^{ic} is in fact a root of unity. Building on this, Section 7 uses further results of Boyer and Zhang [BZ98] for small knots to prove Theorem 1.4.

In Section 8, we use the fact that e^{ic} is a root of unity to prove Theorem 8.2, asserting that if the pillowcase image of the $SU(2)$ character variety consists mainly of line segments of slope $-r \in \mathbb{Q}$, then K must be $SU(2)$ -averse. This allows us to prove in Section 9 that $SU(2)$ -averse knots have infinitely many instanton L-space surgeries, and our applications include Theorem 9.3, which says that smoothly slice knots are not $SU(2)$ -averse.

Finally, in Section 10 we investigate when a satellite of a given knot can be $SU(2)$ -averse. We prove the following, a combination of Propositions 10.2 and 10.3:

Theorem 1.7. *Let K be a nontrivial knot, and suppose that some satellite $P(K)$ with winding number w is $SU(2)$ -averse.*

- *If $P(U)$ is not the unknot, then it is also $SU(2)$ -averse, and $r(P(K)) = r(P(U))$.*
- *If $w \neq 0$, then K is $SU(2)$ -averse, with $r(P(K)) = w^2 r(K)$.*

In particular, we prove that $SU(2)$ -averse knots are prime (Theorem 10.4) and we completely determine when a cable of a given knot is $SU(2)$ -averse (Theorem 10.6), leading to a proof of Conjecture 1.5 for algebraic knots (Corollary 10.8). Similarly, in Section 11 we apply many of the above results to prove that several other classes of knots (as described following Conjecture 1.5 above) are not $SU(2)$ -averse. In doing so, we make use of the following strengthening of a theorem of Crowell [Cro59, Theorem 6.5].

Proposition 11.2. *If K is a prime alternating knot, then $\det(K) \geq 3c(K) - 8$ unless K is a $(2, 2k + 1)$ torus knot or a twist knot.*

Acknowledgments. We thank Hans Boden, Marc Culler, Cynthia Curtis, Nathan Dunfield, Paul Feehan, and Michael Heusener for helpful conversations. We are also grateful to the Max Planck Institute for Mathematics for its hospitality during a significant portion of this work. The second author is also grateful for support by the SFB ‘Higher invariants’ (funded by the Deutsche Forschungsgemeinschaft (DFG)) at the University of Regensburg, and for support by a Heisenberg fellowship of the DFG.

2. BACKGROUND

To any space Y we can associate its $SU(2)$ representation variety

$$R(Y) = \text{Hom}(\pi_1(Y), SU(2)).$$

We can realize $R(Y)$ as a real algebraic subset of some \mathbb{R}^n , given a finite presentation

$$\pi_1(Y) = \langle g_1, g_2, \dots, g_k \mid w_1, w_2, \dots, w_l \rangle.$$

Namely, we assign to each g_j four real variables a_j, b_j, c_j, d_j satisfying the relation $a_j^2 + b_j^2 + c_j^2 + d_j^2 = 1$, so that we can set

$$\rho(g_j) = \begin{pmatrix} a_j + ib_j & c_j + id_j \\ -c_j + id_j & a_j - ib_j \end{pmatrix}, \quad \rho(g_j^{-1}) = \begin{pmatrix} a_j - ib_j & -c_j - id_j \\ c_j - id_j & a_j + ib_j \end{pmatrix}.$$

Then each 2×2 matrix equation $\rho(w_i) = I$ gives four additional polynomial relations among the various a, b, c, d , and the collection of these relations defines $R(Y) \subset \mathbb{R}^{4k}$ as the zero locus of some collection of polynomials with coefficients in \mathbb{Z} . (Since $\mathbb{Z}[a_1, \dots, d_k]$ is Noetherian, we only need finitely many of these relations even if $\pi_1(Y)$ was merely finitely generated.) This presents $R(Y)$ as a closed subspace of $(S^3)^k$, hence it is compact.

A representation $\pi_1(Y) \rightarrow SU(2)$ is irreducible if and only if its image does not lie in some $U(1)$ subgroup, or equivalently if its image is not abelian. The space

$$R^*(Y) = \{\rho : \pi_1(Y) \rightarrow SU(2) \mid \rho \text{ irreducible}\}$$

can therefore be cut out from $R(Y)$ by imposing the additional constraint

$$\rho(g_i g_j) \neq \rho(g_j g_i) \text{ for some } i, j.$$

Equivalently, we can sum the squares of the magnitudes of the entries of the matrices $\rho(g_i g_j) - \rho(g_j g_i)$ over all $1 \leq i < j \leq k$, and this is positive if and only if ρ is irreducible, so $R^*(Y)$ is a semi-algebraic set defined by the polynomial relations appearing in $R(Y)$ plus an additional polynomial inequality, again with coefficients in \mathbb{Z} .

Dividing out the action of $SU(2)$ by conjugation on the representation variety, we get the $SU(2)$ character variety and its irreducible part:

$$X(Y) = R(Y)/SU(2), \quad X^*(Y) = R^*(Y)/SU(2).$$

Since $R(Y)$ is compact, so is its quotient $X(Y)$.

In the case of a torus T^2 , we can take a pair of elements μ, λ which generate $\pi_1(T^2) \cong \mathbb{Z}^2$, and since they commute their images under any representation $\rho : \pi_1(T^2) \rightarrow SU(2)$ can be simultaneously diagonalized as

$$\rho(\mu) = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{-i\alpha} \end{pmatrix}, \quad \rho(\lambda) = \begin{pmatrix} e^{i\beta} & 0 \\ 0 & e^{-i\beta} \end{pmatrix}.$$




FIGURE 1. Identifications of the rectangle $[0, \pi] \times [0, 2\pi]$ yielding the pillowcase. The image of $X(K)$ is depicted for K the left-handed trefoil.

The pairs α, β lie in a torus $T = (\mathbb{R}/2\pi\mathbb{Z}) \times (\mathbb{R}/2\pi\mathbb{Z})$, but they are not uniquely determined, since conjugation by $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ induces an involution ι sending (α, β) to $(2\pi - \alpha, 2\pi - \beta)$. Each conjugacy class in $SU(2)$ is completely determined by its trace, so in fact $X(T^2)$ is the quotient of T by this involution:

$$X(T^2) = ((\mathbb{R}/2\pi\mathbb{Z}) \times (\mathbb{R}/2\pi\mathbb{Z}))/\iota. \quad (2.1)$$

The space $X(T^2)$ is often called the *pillowcase*: it is homeomorphic to a sphere with four orbifold points of order 2, since ι has the four fixed points $(0, 0)$, $(0, \pi)$, $(\pi, 0)$, and (π, π) . It can be constructed as the quotient of the fundamental domain $[0, \pi] \times [0, 2\pi]$ by the identifications

$$(0, \beta) \sim (0, 2\pi - \beta), \quad (\alpha, 0) \sim (\alpha, 2\pi), \quad (\pi, \beta) \sim (\pi, 2\pi - \beta)$$

where $0 \leq \alpha \leq \pi$ and $0 \leq \beta \leq 2\pi$, see Figure 1 below. We can cut the pillowcase open along the lines $\alpha = 0$ and $\alpha = \pi$ to get an annulus $[0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z})$, which we will call the *cut-open pillowcase*.

Given a knot $K \subset S^3$, we let $R(K) = R(S^3 \setminus N(K))$ and likewise for $R^*(K)$, $X(K)$, and $X^*(K)$. The inclusion $i : \partial N(K) \hookrightarrow S^3 \setminus N(K)$ of the boundary torus induces a restriction map on the $SU(2)$ character varieties

$$i^* : X(K) \rightarrow X(T^2).$$

The image of $X(K)$ inside the cut-open pillowcase $[0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z})$ has canonical coordinates (α, β) given by taking μ and λ to be a meridian and longitude of the knot. (Explicitly, we have $\alpha = \arccos(\frac{1}{2} \operatorname{tr}(\rho(\mu)))$, where $0 \leq \arccos(\theta) \leq \pi$; and then the uniqueness of β will follow from Proposition 2.1 below.) It is also compact since $X(K)$ is. Figure 1 shows the image of $X(K)$ in the pillowcase in the case where K is the left-handed trefoil.

The following facts are more or less standard; see e.g. [Lin16, Lemma 2.8].

Proposition 2.1. *Let $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ be a representation and (α, β) the coordinates of its image in the pillowcase.*

- (1) *Every point on the line $\beta \equiv 0 \pmod{2\pi}$ is the image of a reducible ρ .*
- (2) *If ρ is reducible, then $\beta \equiv 0 \pmod{2\pi}$.*

- (3) If ρ is reducible and a limit of irreducible representations, then $e^{2i\alpha}$ is a root of the Alexander polynomial $\Delta_K(t)$.
- (4) There is a constant $\varepsilon > 0$ depending only on K such that if ρ is irreducible, then $\varepsilon < \alpha < \pi - \varepsilon$.

Proof. Composing the abelianization $\pi_1(S^3 \setminus K) \rightarrow H_1(S^3 \setminus K) \cong \mathbb{Z}$ with the map $k \mapsto \text{diag}(e^{ik\alpha}, e^{-ik\alpha})$ gives a reducible ρ with coordinates $(\alpha, 0)$, since λ is nullhomologous. Conversely, if ρ is reducible then its image is abelian, so ρ factors through $H_1(S^3 \setminus K)$ and hence λ lies in its kernel. The claim about reducible limits of irreducible representations is a theorem of Klassen [Kla91, Theorem 19].

Finally, if $\alpha = 0$ or $\alpha = \pi$ then $\rho(\mu) = \pm I$, and thus ρ is reducible with image in $\{\pm I\}$ since μ normally generates the knot group. In these cases $\Delta_K(e^{2i\alpha}) = \Delta_K(1) = 1$, so some neighborhood of ρ in $R(K)$ consists only of reducibles. Now if there is a sequence of irreducibles ρ_n with coordinates (α_n, β_n) such that $\alpha_n \rightarrow 0$ (resp. $\alpha_n \rightarrow \pi$), then by compactness some subsequence converges to a representation with α -coordinate 0 (resp. π), which must then have a neighborhood with no irreducibles, giving a contradiction. \square

Remark 2.2. The coordinates on the pillowcase itself are not quite canonical, since $(0, \beta)$ is identified with $(0, 2\pi - \beta)$ and likewise for (π, β) and $(\pi, 2\pi - \beta)$. However, since the image of $X(K)$ avoids all such points except for $(0, 0)$ and $(\pi, 0)$, the coordinates on the image of $X(K)$ are indeed well-defined, both on the pillowcase and on its cut-open variant, and so we can safely ignore this ambiguity.

The pillowcase images of character varieties exhibit symmetries, as in the following two propositions.

Proposition 2.3. *The image of $X(K)$ inside the pillowcase is invariant under the involution $(\alpha, \beta) \mapsto (\pi - \alpha, 2\pi - \beta)$.*

Proof. Suppose that the image of ρ has coordinates (α, β) , and hence $\rho(\mu) = \text{diag}(e^{i\alpha}, e^{-i\alpha})$ and $\rho(\lambda) = \text{diag}(e^{i\beta}, e^{-i\beta})$ up to conjugacy. We take the central character

$$\chi : \pi_1(S^3 \setminus K) \xrightarrow{\text{ab}} \mathbb{Z} \rightarrow \{\pm I\} \subset SU(2)$$

in which the latter map sends $k \in \mathbb{Z}$ to $(-I)^k$, and note that $\chi(\mu) = -I$ and $\chi(\lambda) = I$. The representation $\tilde{\rho} = \chi\rho$ satisfies $\tilde{\rho}(\mu) = \text{diag}(e^{i(\alpha+\pi)}, e^{-i(\alpha+\pi)})$ and $\tilde{\rho}(\lambda) = \text{diag}(e^{i\beta}, e^{-i\beta})$, and since $\iota(\alpha + \pi, \beta) = (\pi - \alpha, 2\pi - \beta)$ lies in $[0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z})$, it follows that the latter are the coordinates of $\tilde{\rho}$ in the pillowcase. \square

Proposition 2.4. *Let \overline{K} denote the mirror of K . Then the pillowcase image of $X(\overline{K})$ is the reflection of the pillowcase image of $X(K)$ across the line $\beta = \pi$, obtained by the transformation $(\alpha, \beta) \mapsto (\alpha, 2\pi - \beta)$.*

Proof. We observe that \overline{K} has the same knot group as K , but with peripheral data

$$\mu_{\overline{K}} = \mu_K^{-1}, \quad \lambda_{\overline{K}} = \lambda_K.$$

Any representation $\rho \in R(K)$ with image (α, β) thus gives rise to an identical $\bar{\rho} \in R(\overline{K})$ with coordinates $\iota(2\pi - \alpha, \beta) = (\alpha, 2\pi - \beta)$. \square

Finally, we have the following general facts about the components of the image of $X(K)$ in the pillowcase.

Proposition 2.5. *The image of $X(K)$ inside the cut-open pillowcase $C = [0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z})$ has finitely many path components. If K is not the unknot, then at least one of these components is homologically nontrivial in $H_1(C; \mathbb{Z}) \cong \mathbb{Z}$.*

Proof. The representation variety $R(K)$ is a semi-algebraic subset of some \mathbb{R}^n , so by [BCR98, Theorem 2.4.5] it has finitely many connected components, each of them semi-algebraic and semi-algebraically connected and therefore path-connected by [BCR98, Proposition 2.5.13]. Its quotient $X(K)$ then has finitely many path components, and so does its image in the pillowcase. (In fact, the image is a finite embedded graph, as argued in [Zen16].) Cutting the pillowcase open along $\alpha = 0$ and $\alpha = \pi$ does not change the number of components, since the image only meets these lines in the points $(0, 0)$ and $(\pi, 0)$.

The existence of a homologically nontrivial component when K is a nontrivial knot is a theorem of the second author [Zen16, Theorem 7.1], which relies in turn on the fact that the result $S_0^3(K)$ of zero-surgery on K separates a symplectic 4-manifold with nonvanishing Donaldson invariants [KM04b, KM10b]. \square

Corollary 2.6. *If K is nontrivial, then the image of $X(K)$ inside the fundamental domain $[0, \pi] \times [0, 2\pi]$ of the cut-open pillowcase contains a path from the line $\beta = 0$ to the line $\beta = 2\pi$.*

Proof. Let $Z = i^*(X(K)) \subset X(T^2)$ denote the image of $X(K)$, and let $z_0 \in Z$ be the image of some reducible representation. If $j : Z \hookrightarrow [0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z})$ denotes inclusion into the cut-open pillowcase and $\pi : [0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z}) \rightarrow \mathbb{R}/2\pi\mathbb{Z}$ is projection, then the composition

$$\pi_1(Z, z_0) \xrightarrow{j^*} \pi_1([0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z}), z_0) \xrightarrow{\pi^*} \pi_1(\mathbb{R}/2\pi\mathbb{Z}, 0)$$

is nonzero, because the abelianization $H_1(Z) \rightarrow H_1([0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z}))$ of the first map is nonzero and the second map is an isomorphism. But $\pi \circ j$ sends a point to its β -coordinate, so if $\gamma \in \pi_1(Z, z_0)$ is a path with image $2\pi n > 0$ in $\pi_1(\mathbb{R}/2\pi\mathbb{Z}, 0) \cong 2\pi\mathbb{Z}$ then we can lift its β -coordinate to \mathbb{R} to get a path from 0 to $2\pi n$, and a segment of this path from $\beta = 0$ to $\beta = 2\pi$ corresponds to the desired path in the fundamental domain. \square

3. THE PILLOWCASE AND $SU(2)$ -CYCLIC SURGERIES

In this section we use the image of $X(K)$ inside the pillowcase to understand which surgeries on K do not admit irreducible $SU(2)$ representations. The fundamental group of $\frac{m}{n}$ -surgery on K is the quotient $\pi_1(S^3 \setminus K) / \langle \mu^m \lambda^n \rangle$, so irreducible representations of the surgered manifold correspond bijectively to irreducible representations ρ of the knot group satisfying $\rho(\mu^m \lambda^n) = I$.

In this section we will let $\pi : \mathbb{R}^2 \rightarrow X(T^2)$ denote the map

$$\mathbb{R}^2 \rightarrow (\mathbb{R}/2\pi\mathbb{Z})^2 \rightarrow X(T^2),$$

defined by as the composition of reduction mod 2π with the quotient by the involution ι of equation (2.1). It is clear that paths in $X(T^2)$ lift via π to paths in \mathbb{R}^2 .

Proposition 3.1. *Let $\gamma : [0, 1] \rightarrow X(T^2)$ be a path consisting of images of irreducible characters of $\pi_1(S^3 \setminus K)$ in the pillowcase. Choose a lift $\tilde{\gamma} : [0, 1] \rightarrow \mathbb{R}^2$ of γ to the plane, so that $\pi \circ \tilde{\gamma} = \gamma$. Suppose that $\tilde{\gamma}(0)$ and $\tilde{\gamma}(1)$ are distinct points in \mathbb{R}^2 with coordinates (α_0, β_0) and (α_1, β_1) , and that some $\frac{m}{n}$ -surgery on K is $SU(2)$ -cyclic with $n \geq 1$.*

- If $\alpha_0 \neq \alpha_1$ and if $r = \frac{\beta_1 - \beta_0}{\alpha_1 - \alpha_0}$ is the slope of the line segment from $\gamma(0)$ to $\gamma(1)$, then

$$\left| \frac{m}{n} - (-r) \right| < \frac{c_\gamma}{n}$$

where $c_\gamma = 2\pi/|\alpha_1 - \alpha_0|$.

- If $\alpha_0 = \alpha_1$, then $n < \frac{2\pi}{|\beta_1 - \beta_0|}$.

Proof. Suppose that each $\tilde{\gamma}(t)$ has coordinates $(\alpha_t, \beta_t) \in \mathbb{R}^2$. If $\frac{m}{n}$ -surgery on K is $SU(2)$ -cyclic, then any representation $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ such that $\rho(\mu^m \lambda^n) = I$ must be reducible. In particular, if ρ is irreducible with image $\tilde{\gamma}(t)$, then $\rho(\mu^m \lambda^n) \neq I$. But up to conjugacy we have $\rho(\mu) = \text{diag}(e^{i\alpha_t}, e^{-i\alpha_t})$ and $\rho(\lambda) = \text{diag}(e^{i\beta_t}, e^{-i\beta_t})$, so since $S^3_{m/n}(K)$ is $SU(2)$ -cyclic we must have

$$m\alpha_t + n\beta_t \not\equiv 0 \pmod{2\pi}$$

for all t , $0 \leq t \leq 1$.

At $t = 0$, the fact that $S^3_{m/n}(K)$ is $SU(2)$ -cyclic tells us that

$$2\pi k < m\alpha_0 + n\beta_0 < 2\pi(k+1)$$

for some integer k . Now $m\alpha_t + n\beta_t$ varies continuously with t , and it can never equal $2\pi k$ or $2\pi(k+1)$ since these are $0 \pmod{2\pi}$, so at time $t = 1$ we must have

$$2\pi k < m\alpha_1 + n\beta_1 < 2\pi(k+1)$$

as well. Combining these inequalities, we see that

$$|(m\alpha_1 + n\beta_1) - (m\alpha_0 + n\beta_0)| < 2\pi.$$

If $\alpha_0 \neq \alpha_1$, then dividing both sides by $n|\alpha_1 - \alpha_0|$ yields

$$\left| \frac{m}{n} + \left(\frac{\beta_1 - \beta_0}{\alpha_1 - \alpha_0} \right) \right| < \frac{2\pi}{n|\alpha_1 - \alpha_0|}.$$

Otherwise $\alpha_0 = \alpha_1$ but $\beta_0 \neq \beta_1$ since $\gamma(0) \neq \gamma(1)$, and we have $n < 2\pi/|\beta_1 - \beta_0|$ as claimed. \square

Theorem 3.2. *Suppose that K admits infinitely many $SU(2)$ -cyclic surgeries. Define*

$$I(K) \subset \mathbb{R}^2$$

as the preimage under $\pi : \mathbb{R}^2 \rightarrow X(T^2)$ of the pillowcase image of $X^(K)$. Then every path component of $I(K)$ is either a point or a line segment, and the line segments all have the same slope. If in addition K is not the unknot, then at least one path component is a line segment, and its slope $r \in \mathbb{R} \cup \{\infty\}$ satisfies $|r| > 2$. If $r \neq \infty$, then the $SU(2)$ -cyclic slopes $\frac{m}{n}$ all satisfy*

$$\left| \frac{m}{n} - (-r) \right| \leq \frac{|r|}{n},$$

so only finitely many values of m are possible for any given n , and $\frac{m}{n} \rightarrow -r$ as $n \rightarrow \infty$.

Proof. Let $I_0 \subset I(K)$ be a path component. If I_0 is not contained in some straight line, then it contains three points $p_i = (\alpha_i, \beta_i)$, $i = 0, 1, 2$, which are not collinear. The α_i cannot all be the same, so at most two of them are equal; we label the p_i so that α_0 is different from both α_1 and α_2 . Letting $c_i = \frac{2\pi}{|\alpha_i - \alpha_0|}$ and $r_i = \frac{\beta_i - \beta_0}{\alpha_i - \alpha_0}$ for $i = 1, 2$, we know that $r_1 \neq r_2$ or else the p_i would be collinear, and that $r_1, r_2 \neq \infty$.

If $\frac{m}{n}$ is an $SU(2)$ -cyclic surgery slope for K , say with $n > 0$, then we can apply Proposition 3.1 to paths from p_0 to p_1 and from p_0 to p_2 within I_0 to see that

$$\left| \frac{m}{n} - (-r_1) \right| < \frac{c_1}{n}, \quad \left| \frac{m}{n} - (-r_2) \right| < \frac{c_2}{n}.$$

Given a sequence of such slopes $\frac{m_k}{n_k}$ with $n_k \rightarrow \infty$, it would follow that $\frac{m_k}{n_k}$ converges to both $-r_1$ and $-r_2$, which contradicts $r_1 \neq r_2$. The $SU(2)$ -cyclic slopes $\frac{m}{n}$ therefore satisfy some uniform upper bound $n \leq C$. But from either of these inequalities it follows that for fixed n there can only be finitely many m such that $\frac{m}{n}$ is an $SU(2)$ -cyclic surgery slope, so the total number of such slopes is finite, which is a contradiction. We conclude that I_0 must be contained in a straight line.

In fact, the above argument shows that the slope r of the straight line is uniquely determined by the set $\{\frac{m_k}{n_k}\}$ of $SU(2)$ -cyclic slopes. Indeed, Proposition 3.1 shows that $r = \infty$ if and only if the n_k are bounded, and if they are unbounded then $n_k \rightarrow \infty$ and $-r = \lim_{n_k \rightarrow \infty} \frac{m_k}{n_k}$. Thus all nontrivial path components of $I(K)$ have the same slope.

If K is not the unknot, then Corollary 2.6 says that there is a path in

$$I(K) \cup (\mathbb{R} \times 2\pi\mathbb{Z}) \subset \mathbb{R}^2,$$

which is the lift to \mathbb{R}^2 of the image of all of $X(K)$ (including the reducible characters), between two points $(\alpha_0, 0)$ and $(\alpha_1, 2\pi)$ with $0 < \alpha_0, \alpha_1 < \pi$. The path must be a straight line in the region $0 < \beta < 2\pi$, where every point is a lift of the image of an irreducible character, and for any small $\delta > 0$ it connects points (α'_0, δ) and $(\alpha'_1, 2\pi - \delta)$ which project to the image of two irreducibles, with slope $r = \frac{2\pi - 2\delta}{\alpha'_1 - \alpha'_0}$.

By Proposition 2.1, there is some $\varepsilon > 0$ depending only on K such that every irreducible character has α -coordinate in the open interval $(\varepsilon, \pi - \varepsilon)$ in the pillowcase. The line segment in \mathbb{R}^2 between (α'_0, δ) and $(\alpha'_1, 2\pi - \delta)$ consists entirely of lifts of images of characters which have $\beta \notin 2\pi\mathbb{Z}$ and are thus irreducible, so it follows that $k\pi + \varepsilon < \alpha'_0, \alpha'_1 < (k+1)\pi - \varepsilon$ for some integer k . This gives us $|\alpha'_1 - \alpha'_0| < \pi - 2\varepsilon$, so the slope r of this segment satisfies $|r| > \frac{2\pi - 2\delta}{\pi - 2\varepsilon}$. Taking limits as $\delta \rightarrow 0$, we see that $|r| \geq \frac{2\pi}{\pi - 2\varepsilon} > 2$. We remark that if $r \neq \infty$ then we also have $|\frac{m}{n} - (-r)| < \frac{2\pi/|\alpha'_1 - \alpha'_0|}{n}$ for each such (α'_0, δ) and $(\alpha'_1, 2\pi - \delta)$, and as $\delta \rightarrow 0$ the ratio $\frac{2\pi}{\alpha'_1 - \alpha'_0}$ approaches the slope $-r$, so that $|\frac{m}{n} - (-r)| \leq \frac{|r|}{n}$. \square

Definition 3.3. *Suppose that $K \subset S^3$ is a nontrivial knot with infinitely many $SU(2)$ -cyclic surgeries. Then we will say that K is $SU(2)$ -averse, and we define the limit slope $r(K) \in \mathbb{R} \cup \{\infty\}$ to be $-r$, where r is the common slope of the line segments in $I(K)$ whose existence is guaranteed by Theorem 3.2.*

Rephrasing Theorem 3.2 in terms of Definition 3.3, we see that an $SU(2)$ -averse knot with limit slope $r(K)$ satisfies $|r(K)| > 2$, and if $r(K) \neq \infty$ then every $SU(2)$ -cyclic slope $\frac{m}{n}$ for K satisfies

$$\left| \frac{m}{n} - r(K) \right| \leq \frac{|r(K)|}{n}.$$

In particular, there are only finitely many possible m for a given n , so the denominators n are unbounded and we have $\frac{m}{n} \rightarrow r(K)$ as $n \rightarrow \infty$. (We will prove in Section 4 that $r(K) \neq \infty$ always holds.)




FIGURE 2. If the image $i^*(X(K))$ contains a component which is curved, as at left, then there are at most finitely many $SU(2)$ -cyclic surgeries by Theorem 3.2. If there are only vertical segments, as at right, then there could be an unbounded set of $SU(2)$ -cyclic surgeries, but we will see in Section 4 that this cannot occur.

Example 3.4. The torus knots $T_{p,q}$ are $SU(2)$ -averse with limit slope $r = pq$, since Moser [Mos71] showed that $(pq + \frac{1}{n})$ -surgery on $T_{p,q}$ is a lens space, and thus $SU(2)$ -cyclic, for all integers $n \neq 0$.

Theorem 3.5. *If K is a nontrivial $SU(2)$ -averse knot with limit slope $r = r(K)$, and if $r < \infty$ (resp. $r = \infty$), then there are finitely many constants c_1, \dots, c_n such that every irreducible $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ has image on one of the lines $\beta = -r\alpha + c_i$ (resp. $\alpha = c_i$) in the cut-open pillowcase.*

Proof. We lift the pillowcase image of $X^*(K)$ to $I(K) \subset \mathbb{R}^2$ as in Theorem 3.2, which guarantees that $I(K)$ is a union of isolated points and line segments of slope $-r$. Then its image I' in the cut-open pillowcase $C \cong [0, \pi] \times (\mathbb{R}/2\pi\mathbb{Z})$, obtained from $I(K) \cap ([0, \pi] \times \mathbb{R})$ by reducing the second coordinate mod 2π , is also a union of such points and line segments.

The set $I' \subset C$ can be identified with the pillowcase image of $X^*(K)$, since the latter avoids a neighborhood of the lines $\alpha = 0$ and $\alpha = \pi$ in $X(T^2)$. Since $R^*(K)$ is a semi-algebraic set, it has only finitely many path components, exactly as argued in the proof of Proposition 2.5, and hence so do its quotient $X^*(K)$ and the image $I' \subset C$ of this quotient. Since each of these components of I' is contained in a line of the form $\beta = -r\alpha + c_i$ (resp. $\alpha = c_i$), we conclude that finitely many c_i suffice. \square

4. FINITENESS OF LIMIT SLOPES

In this section we will show that $SU(2)$ -averse knots have limit slope $r(K) \neq \infty$. The result is a straightforward consequence of the following theorem about the $SU(2)$ character variety of a nontrivial knot.

Theorem 4.1. *Let K be a nontrivial knot in S^3 . Then at least one of the following holds:*

(1) *The pillowcase image of $X^*(K)$ contains the entire curve $\alpha = \frac{\pi}{2}$, or equivalently for every $\beta \in \mathbb{R}/2\pi\mathbb{Z}$ there is an irreducible $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ such that*

$$\rho(\mu) = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad \text{and} \quad \rho(\lambda) = \begin{pmatrix} e^{i\beta} & 0 \\ 0 & e^{-i\beta} \end{pmatrix};$$

(2) *There is a constant $\varepsilon > 0$ and an arc of irreducible representations*

$$\rho_s : \pi_1(S^3 \setminus K) \rightarrow SU(2), \quad s \in \left[\frac{\pi}{2}, \frac{\pi}{2} + \varepsilon \right)$$

with α -coordinate s , i.e. such that $\rho_s(\mu) = \text{diag}(e^{is}, e^{-is})$.

If case (1) applies, then K has no nontrivial $SU(2)$ -cyclic surgeries.

The last claim is immediate: if $\frac{p}{q} \neq \infty$ is an $SU(2)$ -cyclic slope, then the line $p\alpha + q\beta \equiv 0 \pmod{2\pi}$ in the pillowcase intersects the line $\alpha = \frac{\pi}{2}$ at the point

$$\left(\frac{\pi}{2}, -\frac{p\pi}{2q} \pmod{2\pi} \right),$$

so there is no irreducible representation $\pi_1(S^3 \setminus K) \rightarrow SU(2)$ with image at this point. Before proving Theorem 4.1, we explain how the finiteness of $r(K)$ follows.

Theorem 4.2. *If $K \subset S^3$ is a nontrivial $SU(2)$ -averse knot, then the limit slope $r(K)$ is finite.*

Proof. Since K has $SU(2)$ -cyclic surgeries, it must satisfy case (2) of Theorem 4.1. Thus we have an arc of irreducible representations

$$\rho_s : \pi_1(S^3 \setminus K) \rightarrow SU(2), \quad s \in \left[\frac{\pi}{2}, \frac{\pi}{2} + \varepsilon \right)$$

where each ρ_s has α -coordinate s in the pillowcase. If $r(K) = \infty$, then by Theorem 3.5 we can only achieve finitely many α -coordinates along this arc, and this is a contradiction. \square

The proof of Theorem 4.1 follows the same lines as the following theorem of Kronheimer and Mrowka, so we briefly recall its proof here.

Theorem 4.3 ([KM10b, Corollary 7.17]). *Every nontrivial knot $K \subset S^3$ admits a non-abelian representation*

$$\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$$

such that $\rho(\mu)$ is traceless.

Proof (sketch). Kronheimer and Mrowka construct for each knot $K \subset S^3$ a closed 3-manifold

$$Y_1(K) = (S^3 \setminus N(K)) \cup (F \times S^1), \quad (4.1)$$

where F is a genus-1 surface with one boundary component, and the gluing identifies

$$\mu \sim \{p\} \times S^1, \quad \lambda \sim \partial F \times \{q\}.$$

Letting $\alpha, \beta \subset F$ be a pair of curves which intersect transversely in a point, they define a Hermitian line bundle $w \rightarrow Y_1(K)$ such that $c_1(w)$ is Poincaré dual to $\beta \times \{p\} \subset F \times S^1$, and a $U(2)$ -bundle $E \rightarrow Y_1(K)$ equipped with an isomorphism $\wedge^2 E \rightarrow w$. We denote by

$$X^w(Y_1(K)) = \{\rho \in \text{Hom}(\pi_1(Y_1(K)), SO(3)) \mid w_2(\rho) \equiv c_1(w) \pmod{2}\} / SO(3) \quad (4.2)$$

the variety of characters of representations $\pi_1(Y_1(K)) \rightarrow SO(3)$ whose second Stiefel-Whitney class agrees with $c_1(w)$ modulo 2. In [KM10b, Lemma 7.15], they show that $\mathcal{R}^w(Y_1(K))$ can be identified with a double cover of the variety

$$R(K, i) := \left\{ \rho : \pi_1(S^3 \setminus K) \rightarrow SU(2) \mid \rho(\mu) = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \right\}. \quad (4.3)$$

The centralizer of $\text{diag}(i, -i)$ is a $U(1)$ acting on $R(K, i)$ by conjugation; the orbits are a point for the unique reducible $\rho \in R(K, i)$ and S^1 for all irreducible ρ .

Kronheimer and Mrowka define $KHI(K)$ to be a group whose dimension is half that of the instanton homology $I_*(Y_1(K))_w$, which is defined as in [Don02]. If $R(K, i)$ contains only the reducible representation, then the Chern-Simons functional used to construct $I_*(Y_1(K))_w$ has only two critical points, both nondegenerate, and it follows that $\dim(KHI(K)) \leq 1$. But they prove that $\dim(KHI(K)) \geq 2$ whenever K is knotted, and thus $R(K, i)$ must contain at least one irreducible. \square

We will use $KHI(K)$ in a similar way to prove Theorem 4.1. Assuming that the theorem is false for some knot K , we will describe the pillowcase image of $X(K)$ near the line $\alpha = \frac{\pi}{2}$ in Lemma 4.7. Using a generalization of the results of [Zen16], we then perturb the Chern-Simons functional defining $I_*(Y_1(K))_w$ so that its critical point set corresponds not to the curve $\alpha = \frac{\pi}{2}$ (i.e., where $\rho(\mu)$ is conjugate to $\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$), which was used to prove Theorem 4.3, but to some nearby curve in the pillowcase which avoids the image of $X^*(K)$ completely. This would imply that $KHI(K)$ has rank 1 by the same argument (which we generalize in Theorem 4.8), and so K must be the unknot. We will now begin the proof of Theorem 4.1 in earnest.

We denote by \mathcal{A} the space of $SO(3)$ connections on the adjoint bundle $\text{ad}(E)$, or equivalently, the space of $U(2)$ -connections on E which induce some fixed connection in the determinant line bundle w which we suppress from our notation. The character variety $X^w(Y_1(K))$ can be identified with the space of flat $SO(3)$ connections on $\text{ad}(E)$ modulo determinant-1 gauge transformations,

$$X^w(Y_1(K)) \cong \{A \in \mathcal{A} \mid F_A = 0\} / \text{Aut}(E).$$

Equivalently, it can be identified with projectively flat $U(2)$ -connections in E ; this is independent of the choice of fixed connection in the determinant line bundle.

The proof of Theorem 4.1 will make use of a variation of Kronheimer and Mrowka's argument sketched above by applying holonomy perturbations. In order to use these, we set up the notation and recall the main technical results about holonomy perturbations of [Zen16], and we refer to the latter reference for more details on these perturbations. We define a manifold diffeomorphic to the original $Y_1(K)$ from equation (4.1) above by

$$Y_1(K) = (S^3 \setminus N(K)) \cup ([0, 1] \times T^2) \cup (F \times S^1),$$

where we have introduced a thickened cylinder $M = [0, 1] \times T^2$ in between the knot complement and $(F \times S^1)$. We declare that the boundary component $\{0\} \times T^2$ is the one glued to $S^3 \setminus N(K)$, mapping the circles $\{0\} \times S^1 \times \{\text{pt}\}$ to meridians of K and the circles $\{0\} \times \{\text{pt}\} \times S^1$ to longitudes. The boundary component $\{1\} \times T^2$ is glued to $F \times S^1$ with the circle $\{1\} \times S^1 \times \{\text{pt}\}$ mapping to the S^1 -factors, and with the circles $\{1\} \times \{\text{pt}\} \times S^1$ mapping to the boundary of F .

Let $\Sigma = [0, 1] \times S^1$ be the 2-dimensional annulus. For $k = 0, \dots, n-1$, let

$$A_k = \begin{pmatrix} a_k & c_k \\ b_k & d_k \end{pmatrix} \in SL(2, \mathbb{Z})$$

be a sequence of matrices. We define a sequence of embeddings $\iota_k: \Sigma \times S^1 \rightarrow M$ by the formula

$$\left(t, \begin{pmatrix} z \\ w \end{pmatrix} \right) \mapsto \left(\frac{k+t}{n}, \begin{pmatrix} a_k & c_k \\ b_k & d_k \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} \right), \quad (4.4)$$

where we understand $S^1 = \mathbb{R}/\mathbb{Z}$ and write $\Sigma \times S^1$ as $[0, 1] \times T^2$, so that the matrix multiplication is understood in the usual sense. Notice that $\text{Im}(\iota_k) = [\frac{k}{n}, \frac{k+1}{n}] \times T^2 \subseteq M$.

Let μ be a 2-form on Σ with support in the interior of Σ and integral 1. The 2-form $\pi_{\Sigma}^* \mu$ on $\Sigma \times S^1$, defined by pulling back μ by the projection onto the first factor, can be pushed forward via ι_k to obtain a 2-form μ_k on M . Notice that μ_k has compact support in the interior of M , so that μ_k extends to $Y_1(K)$.

For $k = 0, \dots, n-1$, let $\chi_k: SU(2) \rightarrow \mathbb{R}$ be a sequence of class functions. These give rise to 2π -periodic even functions $g_k: \mathbb{R} \rightarrow \mathbb{R}$ via the formula

$$\chi_k \left(\begin{pmatrix} e^{it} & 0 \\ 0 & e^{-it} \end{pmatrix} \right) = g_k(t). \quad (4.5)$$

To each class function $\chi: SU(2) \rightarrow \mathbb{R}$ we associate a Lie-algebra valued, $SU(2)$ -equivariant function $\chi': SU(2) \rightarrow \mathfrak{su}(2)$ which is dual, with respect to the trace, to the differential $d\chi: TSU(2) \rightarrow \mathbb{R}$.

In the sequel, we suppose we have chosen a trivialization of our bundle $E \rightarrow Y_1(K)$ over M . In fact, by our choice of the line bundle w , there is no obstruction to such a trivialization over M . For any point $z \in \Sigma$ there is a loop $\iota_k(\{z\} \times S^1)$ passing through it, and there is precisely one such loop through any point in the image of ι_k . For such a point x in the image, we denote by $\text{Hol}_{\iota_k}(A)_x$ the holonomy of A along the loop through this point. This way $\text{Hol}_{\iota_k}(A)$ can be seen as a section of the automorphism bundle $\text{Aut}(E)$ over the image of ι_k . For a class function χ_k we can therefore define

$$\chi'_k(\text{Hol}_{\iota_k}(A)) \mu_k$$

which is a 2-form with values in the bundle $\text{ad}(E)$ over $Y_1(K)$, and with compact support in the interior of the image of ι_k .

We now define a family, parametrized by $t \in [0, 1]$, of perturbations to the flatness equation on $Y_1(K)$.

Definition 4.4. *Let the function $\theta^t: \mathcal{A} \rightarrow \Omega^2(Y_1(K); \text{ad}(E))$ be defined in the following way. For each $k = 0, \dots, n-1$ and $\frac{k}{n} \leq t \leq \frac{k+1}{n}$, we let*

$$\theta_{\{\iota_k, \chi_k\}}^t(A) := n \left(t - \frac{k}{n} \right) \cdot \chi'_k(\text{Hol}_{\iota_k}(A)) \mu_k + \sum_{l=0}^{k-1} \chi'_l(\text{Hol}_{\iota_l}(A)) \mu_l.$$

This function interpolates between $\theta^0 = 0$ and $\theta^1(A) = \sum_{l=0}^{n-1} \chi'_l(\text{Hol}_{\iota_l}(A)) \mu_l$.

The perturbed flatness equation for a connection $A \in \mathcal{A}$ now is

$$F_A = \theta_{\{\iota_k, \chi_k\}}^t(A), \quad (4.6)$$

and we denote by $X_{\{\iota_k, \chi_k\}}^w(t)(Y_1(K))$ the space of connections $A \in \mathcal{A}$ which solve this equation, modulo the group of determinant-1 automorphisms of E .

The parameter t is not to be confused with the coordinate t on the thickened torus $M = [0, 1] \times T^2$, although there is the accidental coincidence that θ^t has support inside $[0, \frac{1}{n}[nt]] \times T^2 \subseteq M$. At each time t , the set $X_{\{\iota_k, \chi_k\}(t)}^w(Y_1(K))$ arises as the critical set modulo gauge transformations of some perturbation of the Chern-Simons functional, which for $t = 1$ we write as

$$CS + \Phi : \mathcal{A} \rightarrow \mathbb{R} \tag{4.7}$$

where $\Phi(A) = \sum_{k=0}^{n-1} \int_{\Sigma_k} \chi_k(\text{Hol}_{\iota_k}(A)) \mu_k(z)$.

Note that equation (4.6) also makes sense for $SU(2)$ -connections defined over the thickened cylinder M . We denote by $X_{\{\iota_k, \chi_k\}(t)}(M)$ the connections which solve this equation modulo bundle automorphisms. The two inclusions of the boundary components $\{0\} \times T^2$ and $\{1\} \times T^2$ into M induce two restriction maps r_0 and r_1 to the pillowcase $X(T^2)$, to which we give coordinates α, β just as before.

The main result (Theorem 4.2) of [Zen16] states that given any isotopy through area-preserving maps ψ_t of the pillowcase $X(T^2)$, fixing the four singular points, and given some $\varepsilon > 0$, there is a finite sequence of embeddings $\{\iota_k\}$ and there are class functions $\{\chi_k\}$ as above, such that the isotopy ψ_t is ε -close in the C^0 -topology to an isotopy $\bar{\phi}_t$ which also fixes the four singular points. The isotopy $\bar{\phi}_t$ is described in the following way.

The lift ϕ_t of $\bar{\phi}_t$ to the branched double cover $\mathbb{R}^2/2\pi\mathbb{Z}^2$ is an isotopy which for $\frac{k}{n} \leq t \leq \frac{k+1}{n}$ is given by

$$\phi_t = \zeta_{n(t-\frac{k}{n})}^{(k)} \circ \zeta_1^{(k-1)} \circ \dots \circ \zeta_1^{(0)}$$

for any $k = 0, \dots, n-1$. Here the maps $\zeta_s^{(k)}$ are defined for $s \in [0, 1]$, and for $k = 0, \dots, n-1$ by the equation

$$\zeta_s^{(k)} = A_k \circ \chi_{f_k}^s \circ A_k^{-1},$$

where the A_k are elements of $SL(2, \mathbb{Z})$ and

$$\chi_{f_k}^s : \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \mapsto \begin{pmatrix} \alpha + s \cdot f_k(\beta) \\ \beta \end{pmatrix} \tag{4.8}$$

are shearing isotopies. The functions f_k are the derivatives of the functions g_k defined in equation (4.5). These isotopies $\zeta_s^{(k)}$ are in fact shearing isotopies, as defined in equation (1.1) in the introduction.

Associated to the perturbation data $\{\iota_k, \chi_k\}$, there is the *perturbed character variety* $X_{\{\iota_k, \chi_k\}(t)}(M)$ for every $t \in [0, 1]$, for which the two restriction maps r_0 and r_1 are related by the commutative diagram

$$\begin{array}{ccc} & X_{\{\iota_k, \chi_k\}(t)}(M) & \\ r_0 \swarrow & & \searrow r_1 \\ X(T^2) & \xrightarrow{\bar{\phi}_t} & X(T^2). \end{array} \tag{4.9}$$

We will need a slight enhancement of the approximation result [Zen16, Theorem 3.3] which is a part of the statement of [Zen16, Theorem 4.2]. In fact, the approximation holds in any C^k -topology for $k \geq 0$. However, in this article we will only need the fact that we can approximate area-preserving isotopies by shearing isotopies in the C^1 -topology.

We defer the slightly technical proof of the following result to Appendix A. We state it in the $\mathbb{Z}/2$ -equivariant case which is needed for our application, but we remark that it also holds in the non-equivariant version.

Theorem 4.5. *Let*

$$\psi : [0, 1] \times T^2 \rightarrow T^2, \quad (t, (x, y)) \mapsto \psi_t(x, y)$$

be a $\mathbb{Z}/2$ -equivariant smooth isotopy through area-preserving maps, which necessarily fixes the four fixed points of the hyperelliptic involution. Let $r \geq 0$ be given. Then for any $\varepsilon > 0$, there is a $\mathbb{Z}/2$ -equivariant map

$$\phi : [0, 1] \times T^2 \rightarrow T^2, \quad (t, (x, y)) \mapsto \phi_t(x, y)$$

which is continuous in t and smooth in (x, y) , such that:

(i) If d denotes the Euclidean metric on T^2 , then we have

$$\sup_{(x, y) \in T^2} d(\psi_t(x, y), \phi_t(x, y)) + \sum_{l=1}^r \left\| D^{(l)}\psi_t - D^{(l)}\phi_t \right\|_{\infty} < \varepsilon$$

for all $t \in [0, 1]$ and all $(x, y) \in T^2$.

(ii) There is a partition $0 = t_0 < t_1 \cdots < t_{n+1} = 1$ and a $\mathbb{Z}/2$ -equivariant shearing isotopy $(\zeta_s^{(i)})_{s \in [0, 1]}$ for each $i = 0, \dots, n$ such that for any $t_i \leq t \leq t_{i+1}$, we have

$$\phi_t = \zeta_{t-t_i}^{(i)} \circ \phi_{t_i}$$

for all $(x, y) \in T^2$.

In other words, the smooth isotopy (ψ_t) can be C^r -approximated by isotopies (ϕ_t) which are piecewise (in the coordinate t) isotopies through shearings.

The following lemma is the analogue of [KM10b, Lemma 7.15] for the perturbed character variety $X_{\{\iota_k, \chi_k\}(t)}^w(Y_1(K))$; we will mostly be interested in the case $t = 1$. We let C be the circle in $X(T^2)$ defined by $\alpha = \pi/2$, and then we define the circle

$$\tilde{C} \subset (\mathbb{R}/2\pi\mathbb{Z})_{\alpha} \times (\mathbb{R}/2\pi\mathbb{Z})_{\beta}$$

as one of two possible lifts of C to the branched double cover $\tilde{X}(T^2) \cong T^2$ of $X(T^2)$: for concreteness, we will take \tilde{C} to be the circle $\alpha = \frac{\pi}{2}$, rather than $\alpha = \frac{3\pi}{2}$. This is analogous to a choice implicitly made in [KM10b, Lemma 7.15]: if $\rho(\mu)$ is diagonal and traceless then it could be either $\text{diag}(i, -i)$ or $\text{diag}(-i, i)$, and the two choices are equivalent since they are related by conjugation but Kronheimer and Mrowka choose the former.

Lemma 4.6. *The character variety $X_{\{\iota_k, \chi_k\}(t)}^w(Y_1(K))$ is a double cover of the perturbed representation variety*

$$R(K \mid (\phi_t)^{-1}(\tilde{C})) := \left\{ \rho \in R(K) \mid \begin{array}{l} \rho(\mu) = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{-i\alpha} \end{pmatrix}, \rho(\lambda) = \begin{pmatrix} e^{i\beta} & 0 \\ 0 & e^{-i\beta} \end{pmatrix} \\ \text{for some } (\alpha, \beta) \in (\phi_t)^{-1}(\tilde{C}). \end{array} \right\}.$$

Proof. Lemma 7.14 of [KM10b] describes the $SO(3)$ -character variety of $F \times S^1$ with the second Stiefel-Whitney class w as specified above, where the restriction map on characters from $F \times S^1$ to its boundary T^2 identifies these conjugacy classes of representations with the circle which doubly covers the circle $\tilde{C} \subset \tilde{X}(T^2)$. The representations of T^2 from this circle which extend to equivalence classes of perturbed representations of $X_{\{\iota_k, \chi_k\}(t)}^w(Y_1(K))$

on the other side of $F \times S^1$ are precisely those which correspond to points of $(\phi_t)^{-1}(\tilde{C})$ because of the diagram (4.9) from above. \square

Having explained how the above perturbations affect the generators of the KHI chain complex, we now find a convenient choice of curve in $X(T^2)$ to make the generating set as small as possible. The following lemma, which we will use to prove Theorem 4.1, describes the pillowcase image of $X^*(K)$ in a neighborhood of the curve $\alpha = \frac{\pi}{2}$.

Lemma 4.7. *Suppose that $K \subset S^3$ is a nontrivial knot which does not satisfy Theorem 4.1. Then there are constants b and ε , with $0 < \varepsilon < b < \pi - \varepsilon$, such that if (α, β) lies in the pillowcase image of $X^*(K)$ and*

$$\frac{\pi}{2} - \varepsilon < \alpha < \frac{\pi}{2} + \varepsilon,$$

then $\alpha = \frac{\pi}{2}$ and $\beta \notin (b - \varepsilon, b + \varepsilon) \cup (2\pi - b - \varepsilon, 2\pi - b + \varepsilon)$.

Proof. Since case (1) of Theorem 4.1 does not apply, there is some point $(\frac{\pi}{2}, b_0)$ which is not in the pillowcase image of $X^*(K)$. The variety $R(K, i)$ of (4.3) is compact, as it is the closed subset of the compact variety $R(K)$ cut out by the matrix equation $\rho(\mu) = \text{diag}(i, -i)$. Moreover, since $\Delta_K(-1) = \pm \det(K) \neq 0$, Proposition 2.1 says that the unique reducible representation in $R(K, i)$ is not a limit of irreducibles, and so the subset

$$R^*(K, i) \subset R(K, i)$$

consisting of irreducibles is also compact.

The image of the compact set $R^*(K, i)$ in the pillowcase is closed, so it avoids an open neighborhood $U \subset \{\frac{\pi}{2}\} \times \mathbb{R}/2\pi\mathbb{Z}$ of $(\frac{\pi}{2}, b_0)$. Letting $b \notin \{0, \pi\}$ be the β -coordinate of some point in U , we can assume that $0 < b < \pi$ by Proposition 2.3, and we pick $\delta > 0$ so that $0 < b - \delta < b + \delta < \pi$ and the interval

$$\left\{\frac{\pi}{2}\right\} \times (b - \delta, b + \delta)$$

in the pillowcase is disjoint from the image of $X^*(K)$. The interval

$$\left\{\frac{\pi}{2}\right\} \times (2\pi - b - \delta, 2\pi - b + \delta)$$

also avoids the image of $X^*(K)$, again by Proposition 2.3.

Next, since case (2) of Theorem 4.1 is not satisfied, we claim that $\frac{\pi}{2}$ is not a limit point of the set

$$\left\{\alpha \neq \frac{\pi}{2} \mid \exists \rho \in R^*(K) \text{ with pillowcase coordinates } (\alpha, \beta) \text{ for some } \beta\right\}.$$

This will guarantee for some $\varepsilon > 0$ that any irreducible representation with α -coordinate between $\frac{\pi}{2} - \varepsilon$ and $\frac{\pi}{2} + \varepsilon$ actually has α -coordinate $\frac{\pi}{2}$, and taking $\varepsilon < \delta$ without loss of generality, this will complete the proof.

To prove the claim, suppose that there is a sequence of irreducible representations $\rho_n : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ with pillowcase coordinates (α_n, β_n) , such that $\alpha_n \neq \frac{\pi}{2}$ and

$$\lim_{n \rightarrow \infty} \alpha_n = \frac{\pi}{2}.$$

Using Proposition 2.3, we can assume that $\alpha_n > \frac{\pi}{2}$ for all n ; since $R(K)$ is compact and has finitely many path components, we can pass to a subsequence to ensure that they all lie in the same path component C and converge to a limit ρ , with pillowcase coordinates $(\frac{\pi}{2}, \beta)$ for some β . Then ρ is irreducible as argued above, and since C is closed it must also

contain ρ . A continuous semi-algebraic path $\gamma = \gamma_t : [0, 1] \rightarrow C$ from ρ to ρ_1 (which exists by [BCR98, Proposition 2.5.13]) must eventually leave the line $\alpha = \frac{\pi}{2}$ in the pillowcase, since $\alpha_1 > \frac{\pi}{2}$. If t_0 is the last time at which γ_{t_0} meets this line, then for some $\tau > 0$ the arc $\gamma|_{[t_0, t_0 + \tau]}$ consists of points with distinct α -coordinates; note that here we use the fact that γ is not just continuous but semi-algebraic, so that $\cos(\alpha_t) = \frac{1}{2} \operatorname{tr}(\gamma_t(\mu))$ defines a semi-algebraic map $[0, 1] \rightarrow \mathbb{R}$. A reparametrization of this arc shows that K must have satisfied case (2) of Theorem 4.1 after all, which is a contradiction. \square

Lemmas 4.6 and 4.7 now point us toward the proof of Theorem 4.1: we find an isotopy which takes C to some curve in the pillowcase which avoids the image $i^*(X^*(K))$ of the irreducible characters of K , so that $X_{\{\iota_k, \chi_k\}(t)}^w(Y_1(K))$ consists of only finitely many points. If these are irreducible and nondegenerate, then they will generate a complex whose homology is $KHI(K)^{\oplus 2}$. The following statement is more general than we need for Theorem 4.1, but we will eventually use it again in a different setting to prove Theorem 7.3.

Theorem 4.8. *Let $K \subset S^3$ be a knot. Let C' be a smooth, simple closed curve in the pillowcase, and suppose that there is an area-preserving isotopy*

$$\bar{\psi}_t : X(T^2) \rightarrow X(T^2), \quad 0 \leq t \leq 1$$

which takes C' to $C = \{\alpha = \frac{\pi}{2}\}$ and fixes the four orbifold points $(0, 0)$, $(0, \pi)$, $(\pi, 0)$, and (π, π) . Suppose that C' intersects the line $\{\beta \equiv 0 \pmod{2\pi}\}$ transversely in exactly $n \geq 1$ points $(\alpha_1, 0), \dots, (\alpha_n, 0)$, and that $e^{2i\alpha_j}$ is not a root of $\Delta_K(t)$ for any j . If the pillowcase image $i^(X^*(K))$ of the irreducible character variety of K is disjoint from C' , then $KHI(K)$ has rank at most n .*

Proof. We first claim that C' has an open neighborhood U which is disjoint from $i^*(X^*(K))$. Supposing otherwise, we can find conjugacy classes $[\rho_n] \in X^*(K)$ such that the distance from $i^*([\rho_n])$ to C' goes to zero as $n \rightarrow \infty$, and since $R(K)$ is compact, some subsequence of the ρ_n converges to a representation ρ ; since $i^*([\rho])$ has distance zero from the compact set C' , it must lie in C' . By hypothesis ρ must then be reducible, but it is a limit of irreducibles and so it must have pillowcase coordinates $(\alpha, 0)$ with $\Delta_K(e^{2i\alpha}) = 0$. We have assumed that C' avoids all such points, though, giving a contradiction.

Fixing $\varepsilon > 0$, we now lift $\bar{\psi}_t$ to an isotopy $\psi_t : T^2 \rightarrow T^2$ of the branched double cover, apply Theorem 4.5 to find a composition of $\mathbb{Z}/2$ -equivariant shearing isotopies $\phi_t : T^2 \rightarrow T^2$ such that $\|\phi_t - \psi_t\|_{C^1} < \varepsilon$ for all $t \in [0, 1]$, and project this back down to get an isotopy $\bar{\phi}_t : X(T^2) \rightarrow X(T^2)$ which is C^1 -close to $\bar{\psi}_t$. If we take ε sufficiently small, then this will guarantee first that $(\bar{\phi}_1)^{-1}(C)$ lies in the neighborhood U of $C' = (\bar{\psi}_1)^{-1}(C)$, hence remains disjoint from $i^*(X^*(K))$; second, that it still avoids the points $(\alpha, 0)$ with $\Delta_K(e^{2i\alpha}) = 0$; and third, that it still intersects the line $\{\beta \equiv 0 \pmod{2\pi}\}$ transversely in n points. Thus we may replace C' with $(\bar{\phi}_1)^{-1}(C)$ and replace $\bar{\psi}_t$ with the isotopy $\bar{\phi}_t$ which lifts to ϕ_t .

We now use the shearing isotopies which make up ϕ_t to perturb the flatness equation on

$$Y_1(K) = (S^3 \setminus N(K)) \cup ([0, 1] \times T^2) \cup (F \times S^1)$$

to $F_A = \theta_{\{\iota_k, \chi_k\}}^t(A)$, exactly as described in Definition 4.4. At time $t = 1$, we recall that these perturbed flat connections are equivalently the critical points of a perturbed Chern-Simons functional $CS + \Phi$, as in (4.7). By Lemma 4.6, the space $X_{\{\iota_k, \chi_k\}(1)}^w(Y_1(K))$ of solutions to this equation modulo gauge transformations is a double cover of

$$R(K \mid (\phi_1)^{-1}(\tilde{C})) = R(K \mid \tilde{C}'),$$

where \tilde{C} is the lift of C to T^2 satisfying $\alpha = \frac{\pi}{2}$ and $\tilde{C}' = (\phi_1)^{-1}(\tilde{C})$ is the corresponding lift of C' . Thus by hypothesis $X_{\{\iota_k, \chi_k\}(1)}^w(Y_1(K))$ consists of $2n$ points A_1, \dots, A_{2n} .

Our goal is to associate an instanton Floer homology group $I_*^{\{\iota_k, \chi_k\}(1)}(Y_1(K))_w$ to this perturbation data and show that it is isomorphic to the usual group $I_*(Y_1(K))_w$; then the rank of the latter will be at most $2n$, and hence by definition $KHI(K)$ will have rank at most n . In order to carry this out, we must first show that all points of $X_{\{\iota_k, \chi_k\}(1)}^w(Y_1(K))$ are irreducible and nondegenerate.

For irreducibility, we note that the 2-forms μ_k appearing in the definition of $\theta_{\{\iota_k, \chi_k\}}^t(A)$ are supported entirely on $[0, 1] \times T^2$. In particular, any connection A with equivalence class

$$[A] \in X_{\{\iota_k, \chi_k\}(t)}^w(Y_1(K))$$

restricts to a flat connection on both $S^3 \setminus N(K)$ and $F \times S^1$. But flat connections on the bundle $\text{ad}(E)|_{F \times S^1}$ are irreducible by construction, since $c_1(w)$ has odd pairing with the surface $\alpha \times S^1 \subset F \times S^1$ (where E , w , and α are as described in the proof of Theorem 4.3), and so $X_{\{\iota_k, \chi_k\}(t)}^w(Y_1(K))$ does not contain any reducible connections for any t , $0 \leq t \leq 1$.

Now we claim that each $A = A_j \in X_{\{\iota_k, \chi_k\}(1)}^w(Y_1(K))$, $j = 1, \dots, 2n$, is nondegenerate. Writing $Y = Y_1(K)$, this is equivalent to the vanishing of the kernel of the operator

$$K_{A, \Phi} = \begin{pmatrix} 0 & d_A^* \\ d_A & -*d_{A, \Phi} \end{pmatrix}$$

on $(\Omega^0 \oplus \Omega^1)(Y; \text{ad}(E))$, where d_A and d_A^* are the exterior derivative twisted by A and its Hodge dual, and $d_{A, \Phi} = d_A - 4\pi^2 * \text{Hess}_A(\Phi)$; see [Flo88, Tau90] or [Her94, Section 6]. By Hodge theory we can identify

$$\ker(K_{A, \Phi}) \cong H_A^0(Y; \text{ad}(E)) \oplus H_{A, \Phi}^1(Y; \text{ad}(E)),$$

where the groups on the right are cohomology groups of the complex

$$0 \rightarrow \Omega^0(Y; \text{ad}(E)) \xrightarrow{d_A} \Omega^1(Y; \text{ad}(E)) \xrightarrow{d_{A, \Phi}} \Omega^2(Y; \text{ad}(E)) \xrightarrow{d_A} \Omega^3(Y; \text{ad}(E)) \rightarrow 0.$$

Since A is irreducible we have $H_A^0(Y; \text{ad}(E)) = 0$, so we wish to prove that $H_{A, \Phi}^1(Y; \text{ad}(E)) = 0$ as well. In the sequel we will drop the $\text{ad}(E)$ from our notation.

This cohomology can be computed by a Mayer-Vietoris argument, in which we write

$$Y = (S^3 \setminus N(K)) \cup_{T^2} V,$$

where V is a neighborhood of $([0, 1] \times T^2) \cup (F \times S^1)$, so that the perturbation Φ is supported in V away from a collar neighborhood of $\partial V \cong T^2$. Since A is a reducible flat connection on both $S^3 \setminus N(K)$ and T^2 with the same $U(1)$ stabilizer, the restriction map

$$H_A^0(S^3 \setminus N(K)) \rightarrow H_A^0(T^2)$$

is surjective. Thus we have an exact sequence

$$0 \rightarrow H_{A, \Phi}^1(Y) \rightarrow H_A^1(S^3 \setminus N(K)) \oplus H_{A, \Phi}^1(V) \xrightarrow{h} H_A^1(T^2), \quad (4.10)$$

and it suffices to show that the map h is injective. (We may drop Φ from two of the above subscripts because it is zero on $S^3 \setminus N(K)$ and on T^2 .)

Denoting by $\rho = \text{Hol}(A)$ the holonomy representation of A , with restrictions ρ_T and ρ_K to T^2 and $S^3 \setminus N(K)$ having centralizers $Z(\rho_T)$ and $Z(\rho_K)$ respectively, we know that

$$\dim T_{\rho_T} R(T^2) = 3 + \dim Z(\rho_T) = 4, \quad \dim T_{\rho_K} R(K) = 3,$$

by [Gol84, Section 1.2] and [Kla91, Theorem 19] respectively; the latter claim uses the assertion that $\Delta_K(e^{2i\alpha}) \neq 0$. Then we have

$$\dim H_A^1(T^2) = \dim T_{\rho_T} R(T^2) - (3 - \dim Z(\rho_T)),$$

since the tangent spaces at ρ_T to $R(T^2)$ and to the $SU(2)$ -orbit of ρ_T are identified with $\ker(d_A|_{\Omega^1(T^2)})$ and $\text{Im}(d_A|_{\Omega^0(T^2)})$ respectively, and likewise for ρ_K . By construction we have $\dim Z(\rho_T) = \dim Z(\rho_K) = 1$, since ρ_T and ρ_K are reducible (with image in a $U(1)$ subgroup of $SU(2)$) but not central. Thus we have

$$\dim H_A^1(T^2) = 2, \quad \dim H_A^1(S^3 \setminus N(K)) = 1.$$

We will show that $\dim(H_{A,\Phi}^1(V)) = 1$. In this case, since each H^1 is identified with the tangent space to the corresponding moduli space at $[A]$, the assumption that C' meets the line $\beta \equiv 0 \pmod{2\pi}$ transversely at the image of $[\rho] = [\text{Hol}(A)]$ will guarantee that the map h in (4.10) is injective, and hence that $H_{A,\Phi}^1(Y) = 0$.

We decompose $V = ([0, 1] \times T^2) \cup_{T^2} (F \times S^1)$, where we have slightly enlarged the interval $[0, 1]$ so that Φ is supported on the interior of $[0, 1] \times T^2$, and where the intersection T^2 is identified with $\{1\} \times T^2 \subset [0, 1] \times T^2$. Again the Mayer-Vietoris sequence says that

$$0 \rightarrow H_{A,\Phi}^1(V) \xrightarrow{i} H_{A,\Phi}^1([0, 1] \times T^2) \oplus H_A^1(F \times S^1) \xrightarrow{j} H_A^1(T^2)$$

is exact, since the restriction map $H_A^0([0, 1] \times T^2) \rightarrow H_A^0(T^2)$ is surjective. But restriction of connections from $[0, 1] \times T^2$ to T^2 induces an isomorphism of the respective (perturbed) flat moduli spaces, hence the map $H_{A,\Phi}^1([0, 1] \times T^2) \rightarrow H_A^1(T^2)$ between their tangent spaces at $[A]$ is an isomorphism. It follows that $\ker(j)$ is isomorphic to $H_A^1(F \times S^1)$, which is 1-dimensional, and since $\text{Im}(i) \cong H_{A,\Phi}^1(V)$ we conclude that $\dim H_{A,\Phi}^1(V) = 1$, as desired.

The above argument has shown that each connection in the moduli space

$$X_{\{\ell_k, \chi_k\}(1)}^w(Y) = \{A_1, \dots, A_{2n}\}$$

is irreducible and nondegenerate. Thus by standard methods, the perturbed Chern-Simons functional $CS + \Phi$ gives rise to an instanton Floer homology group $I_*^{\{\ell_k, \chi_k\}(1)}(Y)_w$, and we have shown that its rank is at most $2n$. Since there are no reducible connections in any of the moduli spaces $X_{\{\ell_k, \chi_k\}(t)}^w(Y)$ for $0 \leq t \leq 1$, the cobordism argument presented in [Don02, Section 5.3] carries over to our situation and shows that $I_*^{\{\ell_k, \chi_k\}(1)}(Y)_w \cong I_*(Y)_w$. Then $KHI(K)$ is by definition the $+2$ -eigenspace of the operator $\mu(\text{pt}) : I_*(Y)_w \rightarrow I_*(Y)_w$, which is isomorphic to the -2 -eigenspace of $\mu(\text{pt})$ and hence has rank at most n . \square

Here we remark on some points arising in the proof of Theorem 4.8. First, we could make the moduli space of perturbed flat connections regular by a generic holonomy perturbation, but such a perturbation cannot necessarily be confined to the $[0, 1] \times T^2$ region of $Y_1(K)$: in general, one may need to introduce holonomy perturbations along a collection of curves which generates $\pi_1(Y_1(K))$ (see [Don02, Section 5.5.1]), and this destroys the relationship of Lemma 4.6 between the perturbed moduli space and the pillowcase image of $R(K)$. Thus we prefer to check directly that the relevant moduli space is already regular.

Second, in constructing instanton homology groups over 3-manifolds equipped with non-trivial bundles, one usually only allows small perturbations of the Chern-Simons functional in order to ensure that all critical points remain irreducible. However, this is not a concern for us since our perturbed flat connections are still flat on $\text{ad}(E)|_{F \times S^1}$ and hence irreducible.




FIGURE 3. The curves Z and $c_0 = \{\alpha = \frac{\pi}{2}\}$ in the pillowcase.

Instanton homology groups defined with large perturbations have occasionally appeared in the literature before, e.g. in Floer’s proof of the surgery exact triangle [BD95].

We can now complete the proof of Theorem 4.1.

Proof of Theorem 4.1. Suppose that K is a non-trivial knot and neither property (1) nor (2) in the statement of Theorem 4.1 holds. By Lemma 4.7, there are some $b \in (0, \pi)$ and $\varepsilon > 0$ such that the line segment

$$L = \left\{ \left(\frac{\pi}{2}, \beta \right) \mid \beta \in (b - \varepsilon, b + \varepsilon) \right\}$$

of the pillowcase does not meet either of the lines $\beta = 0$ and $\beta = \pi$, and it contains no element of the image of $R^*(K, i)$. (The same statement holds for the image of L under the involution $(\alpha, \beta) \mapsto (\pi - \alpha, 2\pi - \beta)$ of Proposition 2.3.) Furthermore, any irreducible character in $X^*(K)$ whose image in the pillowcase $X(T^2)$ has α -coordinate satisfying $\frac{\pi}{2} - \varepsilon < \alpha < \frac{\pi}{2} + \varepsilon$ must actually have α -coordinate equal to $\frac{\pi}{2}$.

We may suppose that ε is smaller than $\pi/2$. Let $\delta = \frac{\varepsilon^2}{4\pi - 2\varepsilon}$. Let $C \subset X(T^2)$ be the embedded circle in the pillowcase defined by $\alpha = \frac{\pi}{2}$, and let $Z \subset X(T^2)$ be the piecewise linear closed curve which passes through the successive points

$$\begin{aligned} z_0 &= \left(\frac{\pi}{2} - \delta, 0 \right), \\ z_1 &= \left(\frac{\pi}{2} - \delta, b - \frac{\varepsilon}{2} \right), \\ z_2 &= \left(\frac{\pi}{2} + \frac{\varepsilon}{2}, b - \frac{\varepsilon}{2} \right), \\ z_3 &= \left(\frac{\pi}{2} + \frac{\varepsilon}{2}, b + \frac{\varepsilon}{2} \right), \\ z_4 &= \left(\frac{\pi}{2} - \delta, b + \frac{\varepsilon}{2} \right), \\ z_5 &= \left(\frac{\pi}{2} - \delta, 2\pi \right) = z_0. \end{aligned}$$

The value of δ is chosen such that the curve Z separates the pillowcase $X(T^2)$ into two halves of equal area with respect to the standard Euclidean area measure. See Figure 3.

It is clear that we can continuously approximate Z arbitrarily well by rounding the corners $z_1, z_2, z_3,$ and z_4 and fixing an open neighborhood of $z_0 = z_5$ to get a new curve C' , which is isotopic to C through an area-preserving isotopy which fixes the corners of the pillowcase. Then C' still intersects the line $\{\beta \equiv 0 \pmod{2\pi}\}$ transversely in the

single point $(\alpha, 0) = (\frac{\pi}{2} - \delta, 0)$, and for ε sufficiently small we must have $\Delta_K(e^{2i\alpha}) \neq 0$, since $\Delta_K(e^{2i\pi/2}) \neq 0$. We now apply Theorem 4.8 to the curve C' to conclude that $\text{rank}(KHI(K)) \leq 1$. But this is a contradiction just as in the proof of Theorem 4.3, since $KHI(K)$ has rank strictly greater than 1 unless K is unknotted. \square

5. RATIONALITY OF LIMIT SLOPES

In this section we prove that limit slopes of $SU(2)$ -averse knots are always rational numbers, having already shown in Theorem 4.2 that they are finite. In what follows we will let $\overline{\mathbb{Q}} \subset \mathbb{C}$ denote the field of algebraic numbers, and $\mathbb{R}^{\text{alg}} = \overline{\mathbb{Q}} \cap \mathbb{R}$ will denote the field of real algebraic numbers.

Proposition 5.1. *Let K be nontrivial and $SU(2)$ -averse with limit slope $r = r(K)$, and let (α_0, β_0) be a point in the pillowcase image of $X^*(K)$.*

- (1) *If $e^{i\alpha_0}$ is algebraic, then $e^{i\beta_0}$ is algebraic.*
- (2) *If $e^{i\beta_0}$ is algebraic, then $e^{i\alpha_0}$ is algebraic.*
- (3) *If (α_0, β_0) is an isolated point in the image of $X^*(K)$, then $e^{i\alpha_0}$ and $e^{i\beta_0}$ are both algebraic numbers.*

Proof. Certainly $e^{i\alpha_0}$ and $e^{i\beta_0}$ are algebraic if and only if $e^{i(\pi-\alpha_0)}$ and $e^{i(2\pi-\beta_0)}$ are, so by Proposition 2.3 we can replace (α_0, β_0) with $(\pi - \alpha_0, 2\pi - \beta_0)$ as needed to ensure that $0 \leq \beta_0 \leq \pi$. We will take $\rho_0 : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ to be an irreducible representation with $\rho_0(\mu) = \text{diag}(e^{i\alpha_0}, e^{-i\alpha_0})$ and $\rho_0(\lambda) = \text{diag}(e^{i\beta_0}, e^{-i\beta_0})$.

Theorem 3.5 implies that any sufficiently small neighborhood of (α_0, β_0) intersects the pillowcase image of $X^*(K)$ only in the point (α_0, β_0) if it is isolated, or in a connected segment ℓ of the line of slope $-r$ through (α_0, β_0) otherwise. Therefore for sufficiently small rational $\varepsilon > 0$, if (α, β) is in the image of $X^*(K)$ and

$$|\cos(\alpha) - \cos(\alpha_0)|^2 + |\cos(\beta) - \cos(\beta_0)|^2 < 2\varepsilon^2,$$

then either $\beta > \pi$, or $(\alpha, \beta) = (\alpha_0, \beta_0)$, or $(\alpha, \beta) \in \ell$. We then pick rational numbers a and b so that

$$|a - 2\cos(\alpha_0)| < \varepsilon \quad \text{and} \quad |b - 2\cos(\beta_0)| < \varepsilon,$$

and we define a semi-algebraic subset R_0 of $R^*(K)$ by imposing the inequalities

$$(\text{tr}(\rho(\mu)) - a)^2 < \varepsilon^2 \quad \text{and} \quad (\text{tr}(\rho(\lambda)) - b)^2 < \varepsilon^2,$$

as well as the relations $\rho(\mu) = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$ and $\rho(\lambda) = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$, and the additional inequalities $\text{Im}(\rho(\mu)_{1,1}) \geq 0$ and $\text{Im}(\rho(\lambda)_{1,1}) \geq 0$. Here $\text{Im}(\rho(g)_{i,j})$ denotes the imaginary part of the (i, j) -entry of the matrix $\rho(g)$; this is always a polynomial with \mathbb{Z} -coefficients in the variables used to construct $R^*(K)$. The relations on $\rho(\mu)$ and $\rho(\lambda)$ require them to have the form $\text{diag}(e^{i\alpha}, e^{-i\alpha})$ and $\text{diag}(e^{i\beta}, e^{-i\beta})$ respectively. Then $\text{Im}(\rho(\mu)_{1,1}) = \sin(\alpha)$ and $\text{Im}(\rho(\lambda)_{1,1}) = \sin(\beta)$, both of which must be nonnegative, so we can take $0 \leq \alpha, \beta \leq \pi$ at any such $\rho \in R_0$, and then ρ has pillowcase coordinates (α, β) .

We will also impose one additional relation on R_0 in cases (1) and (2). If we are in case (1), so that $e^{i\alpha_0}$ is algebraic, we also impose the relation $\text{tr}(\rho(\mu)) = 2\cos(\alpha_0)$ on R_0 . Similarly, in case (2) we impose $\text{tr}(\rho(\lambda)) = 2\cos(\beta_0)$. We note that $2\cos(\alpha_0)$ (resp. $2\cos(\beta_0)$) is algebraic whenever $e^{i\alpha_0}$ (resp. $e^{i\beta_0}$) is.

In each case R_0 is defined over the real closed field \mathbb{R}^{alg} , and \mathbb{R} is a real closed extension of \mathbb{R}^{alg} , so the Tarski-Seidenberg principle implies that R_0 has an \mathbb{R}^{alg} -point if and only if it has an \mathbb{R} -point [BCR98, Proposition 4.1.1]. But R_0 contains the \mathbb{R} -point ρ_0 by construction,

so R_0 must also have an \mathbb{R}^{alg} -point ρ_{alg} , for which the real and imaginary parts of the entries of every matrix $\rho_{\text{alg}}(g)$, $g \in \pi_1(S^3 \setminus K)$, are real algebraic numbers. In particular, if ρ_{alg} has image (α, β) in the pillowcase, then we have

$$\rho_{\text{alg}}(\mu) = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{-i\alpha} \end{pmatrix}, \quad \rho_{\text{alg}}(\lambda) = \begin{pmatrix} e^{i\beta} & 0 \\ 0 & e^{-i\beta} \end{pmatrix}$$

where $0 \leq \alpha, \beta \leq \pi$, and $e^{i\alpha}$ and $e^{i\beta}$ are both algebraic numbers.

Since $\rho_{\text{alg}} \in R_0$, we have $|2 \cos(\alpha) - a| = |\text{tr}(\rho(\mu)) - a| < \varepsilon$ and $|a - 2 \cos(\alpha_0)| < \varepsilon$, so $|\cos(\alpha) - \cos(\alpha_0)| < \varepsilon$ by the triangle inequality; likewise $|\cos(\beta) - \cos(\beta_0)| < \varepsilon$. Since we ensured that $\beta \leq \pi$, it follows that if (α_0, β_0) is an isolated point then we must have $(\alpha, \beta) = (\alpha_0, \beta_0)$, and that otherwise $(\alpha, \beta) \in \ell$. The latter only applies to cases (1) and (2) of the proposition, and then we know that (α_0, β_0) is the only point of ℓ satisfying $\alpha = \alpha_0$ or $\beta = \beta_0$, so again we must have $(\alpha, \beta) = (\alpha_0, \beta_0)$. In each case we conclude that $e^{i\alpha_0}$ and $e^{i\beta_0}$ are algebraic, since this is true of $e^{i\alpha}$ and $e^{i\beta}$. \square

Proposition 5.2. *Let K be a nontrivial SU(2)-averse knot with limit slope $r = r(K)$, and suppose that the pillowcase image of $X(K)$ contains a nontrivial segment of the line $\beta = -r\alpha + c$. Then $e^{i\pi r}$ and e^{ic} are both algebraic numbers.*

Proof. We pass to a fundamental domain $[0, \pi] \times [0, 2\pi]$ for the cut-open pillowcase. Since $r \neq \infty$, we can pick two points (α_1, β_1) and (α_2, β_2) in this domain, with $\alpha_1 < \alpha_2$, such that the image of $X(K)$ contains the entire line segment ℓ connecting these points.

If we fix an integer $n > \frac{2\pi}{\alpha_2 - \alpha_1}$ and let $m = \lceil \frac{\alpha_1 n}{\pi} \rceil$, then ℓ contains two points of the form $p = (\frac{m}{n}\pi, \beta_-)$ and $p' = (\frac{m+1}{n}\pi, \beta_+)$, and its slope is

$$-r = \frac{n(\beta_+ - \beta_-)}{\pi}.$$

Both $e^{im\pi/n}$ and $e^{i(m+1)\pi/n}$ are algebraic numbers, so Proposition 5.1 says that $e^{i\beta_-}$ and $e^{i\beta_+}$ are algebraic as well, and hence so is

$$\left(\frac{e^{i\beta_-}}{e^{i\beta_+}} \right)^n = e^{-in(\beta_+ - \beta_-)} = e^{i\pi r}.$$

Finally, we rearrange $\beta_- = -r(\frac{m}{n}\pi) + c$ and apply $x \mapsto \exp(ix)$ to both sides to get

$$e^{ic} = e^{i\beta_-} \cdot (e^{i\pi r})^{m/n},$$

and since both $e^{i\beta_-}$ and $e^{i\pi r}$ are algebraic it follows that e^{ic} is as well. \square

Remark 5.3. We will eventually show in Theorem 6.3 that the constants e^{ic} appearing in Proposition 5.2 are in fact roots of unity. The proof will require the fact that r is rational, which we are about to prove.

Theorem 5.4. *Let K be a nontrivial SU(2)-averse knot with limit slope $r = r(K)$. Then r is a rational number.*

Proof. As in the proof of Proposition 5.2, we take two points $p = (\frac{m}{n}\pi, \beta_-)$ and $p' = (\frac{m+1}{n}\pi, \beta_+)$ in the fundamental domain $[0, \pi] \times [0, 2\pi]$ of the pillowcase (so that $0 \leq m < n$) such that the entire line segment ℓ from p to p' lies in the pillowcase image of $X^*(K)$. The line containing ℓ has the form $\beta = -r\alpha + c$, and both $e^{i\pi r}$ and e^{ic} are algebraic.

Given any α such that $e^{i\alpha}$ is algebraic and $\frac{m}{n}\pi < \alpha < \frac{m+1}{n}\pi$, we know that ℓ contains a point $(\alpha, \beta) = (\alpha, -r\alpha + c)$ in the image of $X^*(K)$, and that $e^{i\beta}$ is algebraic by Proposition 5.1. But then

$$e^{ir\alpha} = e^{i(c-\beta)} = \frac{e^{ic}}{e^{i\beta}}$$

is also an algebraic number, so we have shown for α in the range $(\frac{m}{n}\pi, \frac{m+1}{n}\pi)$ that if $e^{i\alpha}$ is algebraic then so is $(e^{i\alpha})^r$. In fact, given any $a \in \mathbb{R}$ such that e^{ia} is algebraic, we can let $\alpha = a + q\pi$ for some rational q such that $\frac{m}{n}\pi < \alpha < \frac{m+1}{n}\pi$; then $e^{i\alpha}$ is also algebraic, so it follows from the above that $(e^{ia})^r = e^{iar} \cdot (e^{i\pi r})^{-q}$ is as well.

Finally, if x_1, x_2, x_3 are nonzero, multiplicatively independent algebraic numbers (meaning that if integers n_1, n_2, n_3 are not all zero, then $x_1^{n_1} x_2^{n_2} x_3^{n_3} \neq 1$), and if each x_j^r is algebraic, then r must be rational [Lan66, §II.1, Corollary 1]. According to Proposition 5.5, we can take $x_j = e^{ia_j}$ for some $a_1, a_2, a_3 \in \mathbb{R}$ such that the numbers a_1, a_2, a_3, π are linearly independent over \mathbb{Q} ; then x_j^r is also algebraic, as argued above. The condition $\prod x_j^{n_j} = 1$ is equivalent to $\sum n_j a_j = 2k\pi$ for some integer k , and then $n_1 = n_2 = n_3 = 0$ by the assumed linear independence, so the x_j are algebraically independent and hence r is rational. \square

The following proposition, which completes the proof of Theorem 5.4, is surely known to experts, but we could not find a proof in the literature so we include one for completeness.

Proposition 5.5. *Let V denote the \mathbb{Q} -vector space of all $x \in \mathbb{R}$ such that e^{ix} is an algebraic number. Then V has infinite dimension over \mathbb{Q} .*

Proof. Suppose that V has a finite basis a_1, \dots, a_n , and let $K = \mathbb{Q}(e^{ia_1}, \dots, e^{ia_n})$. Since each e^{ia_j} is algebraic, K is a finite extension of \mathbb{Q} . Any nonzero $x \in V$ can be written as a \mathbb{Q} -linear combination of the a_i , so upon clearing denominators we have integers $q \geq 1$ and c_1, \dots, c_n such that $qx = c_1 a_1 + c_2 a_2 + \dots + c_n a_n$, and then

$$(e^{ix})^q = (e^{ia_1})^{c_1} (e^{ia_2})^{c_2} \dots (e^{ia_n})^{c_n} \in K.$$

So for every algebraic number $z \in \mathbb{C}$ with $|z| = 1$, there is a positive integer q such that z^q has degree at most $[K : \mathbb{Q}]$ over \mathbb{Q} . If for any $d \geq 1$ we can find an algebraic number z such that $|z| = 1$ and every power z^q ($q \geq 1$) has degree at least d , then taking $d > [K : \mathbb{Q}]$ will give a contradiction, proving that $\dim_{\mathbb{Q}}(V) = \infty$.

In order to find such z , we turn to Salem numbers, for which we refer to the survey [Smy15]. These are real algebraic integers $s > 1$ all of whose Galois conjugates satisfy $|z| \leq 1$, and such that at least one conjugate satisfies $|z| = 1$. It follows from these properties that if s has minimal polynomial $p(t) \in \mathbb{Z}[t]$ of degree d , then $p(s) = p(\frac{1}{s}) = 0$ and all other roots of p lie on the unit circle [Smy15, Lemma 1]; and that s^q is also an algebraic integer of degree d for all $q \geq 1$ [Smy15, Lemma 2].

Let s be a Salem number with minimal polynomial $p(t)$ and splitting field L . Letting z be a root of p with $|z| = 1$ and taking $\sigma \in \text{Gal}(L/\mathbb{Q})$ for which $\sigma(s) = z$, we know that $z^q = \sigma(s^q)$ has the same minimal polynomial as s^q , since if $f \in \mathbb{Z}[t]$ is irreducible and $f(s^q) = 0$ then $f(z^q) = f(\sigma(s^q)) = \sigma(f(s^q)) = 0$ as well. We conclude that z^q is algebraic of degree $\deg(p)$ for all integers $q \geq 1$, so the proof will be complete if we can take $\deg(p)$ arbitrarily large, i.e. if there are Salem numbers of arbitrarily large degree.

Given any fixed $c > 0$, the set of Salem numbers $s \leq c$ of degree at most d is finite. This is because if s is a Salem number with minimal polynomial $p(t) = t^d + a_1 t^{d-1} + \dots + a_d \in \mathbb{Z}[t]$, then each coefficient a_k is (up to sign) the sum of all k -fold products of distinct roots of p ;

there are $\binom{d-1}{k-1}$ such products which involve s and $\binom{d-1}{k}$ which only involve the other roots, and since the other roots all have modulus at most 1, we get bounds $|a_k| \leq c \binom{d-1}{k-1} + \binom{d-1}{k}$. But the set of Salem numbers has limit points, including what are called Pisot numbers [Smy15, Section 3.1] (which include $\sqrt{2} + 1$, for concreteness), so a bounded neighborhood of any limit point contains infinitely many Salem numbers and hence Salem numbers of arbitrarily high degree, as desired. \square

6. THE A-POLYNOMIAL OF AN SU(2)-AVERSE KNOT

The A -polynomial of a knot $K \subset S^3$, defined by Cooper, Culler, Gillet, Long, and Shalen [CCG⁺94], is a polynomial which captures the appropriate analogue of the pillowcase image of $X(K)$ for $SL_2(\mathbb{C})$ representations of $\pi_1(S^3 \setminus K)$. Since $SU(2) \subset SL_2(\mathbb{C})$, it is possible to use what we know about $X(K)$ for an $SU(2)$ -averse knot K to say something about its A -polynomial, and by deep results of [CCG⁺94] we can then in turn understand the geometry of the complement of K . Our goals are to show that the limit slope $r(K)$ is always a *boundary slope* for K , meaning that there is an essential surface in the exterior of K which intersects $\partial N(K)$ in parallel curves of slope $r(K)$ – the set of such slopes for any fixed knot is finite [Hat82] – and to study the constants c_j which appear in Theorem 3.5.

We begin by recalling the construction of the A -polynomial, borrowing our notation from [DG04]. We define the $SL_2(\mathbb{C})$ character variety of a manifold M to be the quotient

$$X_{SL_2}(M) = \text{Hom}(\pi_1(M), SL_2(\mathbb{C})) / SL_2(\mathbb{C}),$$

in which $SL_2(\mathbb{C})$ acts by conjugation. Given a knot $K \subset S^3$ and writing $X_{SL_2}(K)$ in place of $X_{SL_2}(S^3 \setminus N(K))$, the inclusion map $i : T^2 = \partial N(K) \hookrightarrow S^3 \setminus N(K)$ induces a restriction map

$$i^* : X_{SL_2}(K) \rightarrow X_{SL_2}(T^2).$$

The part of $X_{SL_2}(T^2)$ consisting of conjugacy classes of diagonal representations ρ has $\mathbb{C}^* \times \mathbb{C}^*$ as a branched double cover, where we take peripheral elements $\mu, \lambda \in \pi_1(T^2)$ and identify a point $(M, L) \in \mathbb{C}^* \times \mathbb{C}^*$ with the conjugacy class of ρ such that

$$\rho(\mu) = \begin{pmatrix} M & 0 \\ 0 & M^{-1} \end{pmatrix}, \quad \rho(\lambda) = \begin{pmatrix} L & 0 \\ 0 & L^{-1} \end{pmatrix}.$$

(The covering map identifies (M, L) with $(\overline{M}^{-1}, L^{-1})$, exactly as in the $SU(2)$ case.)

We let V be the union of the closures $\overline{i^*(X)}$ over all irreducible components $X \subset X_{SL_2}(K)$ such that $i^*(X)$ has complex dimension 1; by [DG04, Lemma 2.1], we know that $\dim_{\mathbb{C}}(i^*(X))$ is always either 0 or 1. We then let \overline{V} be the preimage of V in $\mathbb{C}^* \times \mathbb{C}^*$. Then \overline{V} is a complex algebraic plane curve, so it is defined by a single polynomial in $\mathbb{C}[M, L]$, which is uniquely determined up to a choice of normalization and can even be taken with integer coefficients. The A -polynomial $A_K(M, L) \in \mathbb{Z}[M, L]$ is this defining polynomial.

In what follows we will also make use of the *Newton polygon* $\mathcal{N}_f \subset \mathbb{R}^2$ of a polynomial $f \in \mathbb{C}[M, L]$: this is the convex hull in the (M, L) -plane of the lattice points $(a, b) \in \mathbb{Z}^2$ for which the $M^a L^b$ -coefficient of f is nonzero. Given two polynomials $f, g \in \mathbb{C}[M, L]$, it is easy to check that \mathcal{N}_{fg} is the *Minkowski sum* of \mathcal{N}_f and \mathcal{N}_g , defined as the set of points $p + q$ where $p \in \mathcal{N}_f$ and $q \in \mathcal{N}_g$. This works equally well for Laurent polynomials $f, g \in \mathbb{C}[M^{\pm 1}, L^{\pm 1}]$, and we will generally consider the A -polynomial in this setting.

Lemma 6.1. *If p, q are relatively prime integers which are not both zero and k is a nonzero complex number, then the polynomial $f(M, L) = M^p L^q - k \in \mathbb{C}[M^{\pm 1}, L^{\pm 1}]$ is irreducible.*

Proof. The Newton polygon \mathcal{N}_f is the line segment from $(0, 0)$ to (p, q) , which only intersects \mathbb{Z}^2 at its two endpoints; there are no lattice points on its interior since p and q are relatively prime. If $f = gh$ where g and h are not monomials, then \mathcal{N}_g and \mathcal{N}_h each contain at least two lattice points as well, so their Minkowski sum $\mathcal{N}_{gh} = \mathcal{N}_f$ must have at least three lattice points and this is impossible. \square

Proposition 6.2. *Let K be a nontrivial knot, and suppose that the pillowcase image of $X^*(K)$ contains a nontrivial line segment of slope $-r \in \mathbb{Q}$; we write $r = \frac{p}{q}$, where p and q are relatively prime integers and $q \geq 1$. If this segment belongs to the line $\beta \equiv -r\alpha + c \pmod{2\pi}$, then $M^p L^q - e^{iqc}$ divides $A_K(M, L)$.*

Proof. By hypothesis there are infinitely many $\rho \in R^*(K)$ whose conjugacy classes have distinct images in the pillowcase $X(T^2)$, and which satisfy $\rho(\mu^p \lambda^q) = \text{diag}(e^{iqc}, e^{-iqc})$. We view these classes $[\rho]$ as points of $X_{SL_2}(K)$ via the inclusion $SU(2) \subset SL_2(\mathbb{C})$. Since $X_{SL_2}(K)$ has finitely many irreducible components, some irreducible component $X \subset X_{SL_2}(K)$ contains infinitely many of the conjugacy classes $[\rho]$, whose images under i^* are all distinct. Then $i^*(X)$ cannot be 0-dimensional, so $\dim_{\mathbb{C}}(i^*(X)) = 1$.

Suppose that the preimage of $i^*(X)$ in $\mathbb{C}^* \times \mathbb{C}^*$ has defining polynomial $P(M, L)$, and let $f(M, L) = M^p L^q - e^{iqc}$. Then the equation $f(M, L) = 0$ is satisfied by infinitely many points of the plane curve $P(M, L) = 0$, namely each of the $i^*([\rho])$ described above, and so $f(M, L)$ and $P(M, L)$ have a common factor. This factor must be $f(M, L) = M^p L^q - e^{iqc}$ itself, since Lemma 6.1 says that f is irreducible, and so f divides P . But P divides $A_K(M, L)$ by definition, so $f(M, L)$ does as well. \square

Theorem 6.3. *Let K be a nontrivial knot, and suppose for some $r \in \mathbb{Q}$ that the pillowcase image of $X^*(K)$ contains a nontrivial segment of the line $\beta \equiv -r\alpha + c \pmod{2\pi}$. Then*

- r is the boundary slope of an essential surface in $S^3 \setminus N(K)$; and
- e^{ic} is a root of unity, or equivalently $c \in 2\pi\mathbb{Q}$.

Proof. Write $r = \frac{p}{q}$ as usual. From Proposition 6.2 we know that $f(M, L) = M^p L^q - e^{iqc}$ divides the A -polynomial of K , so we can write $A_K(M, L) = f(M, L) \cdot g(M, L)$ for some nonzero polynomial g . The Newton polygon \mathcal{N}_{A_K} is the Minkowski sum of \mathcal{N}_f and \mathcal{N}_g , and \mathcal{N}_f is a line segment of slope $\frac{q}{p} = \frac{1}{r}$.

Let $h(M, L)$ be the sum of all monomials of $g(M, L)$ which correspond to points of the vertex or side of \mathcal{N}_g on which $-qM + pL$ is maximized. Then \mathcal{N}_h is either a point or a line segment of slope $\frac{q}{p} = \frac{1}{r}$ on the boundary of \mathcal{N}_g , and \mathcal{N}_{A_K} also has a side E of slope $\frac{1}{r}$, namely the Minkowski sum \mathcal{N}_{fh} of \mathcal{N}_f and \mathcal{N}_h .

Now [CCG⁺94, Theorem 3.4] says that the slopes of the sides of \mathcal{N}_{A_K} are all boundary slopes of essential surfaces. Their result is stated in the (L, M) -plane whereas we have described \mathcal{N}_{A_K} in the (M, L) -plane, so the side E of \mathcal{N}_{A_K} with slope $\frac{1}{r}$ actually produces a surface with boundary slope r , as desired.

In fact, if we write $A_K(M, L) = \sum_{m,n} b_{mn} M^m L^n$ and define the *edge polynomial*

$$\Theta_E(z) = \sum_{(m,n) \in E} b_{mn} z^m,$$

then [CCG⁺94, Proposition 5.10] asserts that $\Theta_E(z)$ is a product of cyclotomic polynomials. Writing $h(M, L) = \sum_{j=0}^k a_j M^{pj+m_0} L^{qj+l_0}$ for some integers k, m_0, l_0 and constants a_j , and

recalling that E is the Newton polygon of fh , we have

$$\Theta_E(z) = (z^p - e^{iqc}) \left(\sum_{j=0}^k a_j z^{pj+m_0} \right).$$

In particular, the factor $z^p - e^{iqc}$ divides a product of cyclotomic polynomials, so its roots are all roots of unity. But these roots are rational powers of e^{ic} , and therefore e^{ic} must be a root of unity as well. \square

Theorem 6.4. *Let K be a nontrivial $SU(2)$ -averse knot. Then the limit slope $r = r(K)$ is a boundary slope for K , and the pillowcase image of $X^*(K)$ consists of finitely many isolated points together with finitely many nontrivial segments of lines $\beta \equiv -r\alpha + c_j$, where $c_j \in 2\pi\mathbb{Q}$ for all j .*

Proof. Theorem 3.2 guarantees the existence of a nontrivial segment of slope $-r$ in the image of $X^*(K)$, and $r \in \mathbb{Q}$ by Theorems 4.2 and 5.4. Theorem 6.3 now says that r is a boundary slope, and that for all such segments the corresponding constant c_j is a rational multiple of 2π . \square

Theorem 6.4 gives us a criterion in terms of Alexander polynomials by which we can show that many knots are not $SU(2)$ -averse.

Proposition 6.5. *Let $K \subset S^3$ be a knot with Alexander polynomial $\Delta_K(t)$, normalized so that $\Delta_K(t) = \Delta_K(t^{-1})$ and $\Delta_K(1) = 1$. If $\Delta_K(t)$ has a root $z = e^{i\theta}$ of odd order which is not a root of unity, then K is not $SU(2)$ -averse. In particular, if $\Delta_K(t)$ has no cyclotomic factors and $\Delta_K(t_0) < 0$ for some $t_0 = e^{i\theta_0}$, then K is not $SU(2)$ -averse.*

Proof. We first check that the given conditions on the factorization and values of $\Delta_K(t)$ suffice to produce a root z of the desired form. Since $\Delta_K(t)$ has real coefficients and $\Delta_K(t) = \Delta_K(t^{-1})$, we have

$$\overline{\Delta_K(e^{i\theta})} = \Delta_K(\overline{e^{i\theta}}) = \Delta_K(e^{-i\theta}) = \Delta_K(e^{i\theta})$$

for all real θ , and so $\Delta_K(e^{i\theta}) \in \mathbb{R}$. The continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(\theta) = \Delta_K(e^{i\theta})$ satisfies $f(0) = \Delta_K(1) = 1$ and $f(\theta_0) = \Delta_K(t_0) < 0$ by hypothesis, so the total multiplicity of the zeroes of f in the interval $[0, \theta_0]$ must be odd, and hence some $\theta \in [0, \theta_0]$ is a zero of odd multiplicity. Then $z = e^{i\theta}$ is a zero of $\Delta_K(t)$ with odd multiplicity, and by assumption it is not a root of unity.

In general, given the odd-order root $z = e^{i\theta}$ and letting $\gamma = \frac{\theta}{2}$, it now follows as in work of Heusener and Kroll [HK98] that the abelian representation $\rho_\gamma : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ with $\rho_\gamma(\mu) = \text{diag}(e^{i\gamma}, e^{-i\gamma})$ is a limit of irreducible representations. In fact, one can adapt their proof of [HK98, Proposition 3.8] to show that the pillowcase image of $X^*(K)$ has $(\gamma, 0)$ as a limit point. The rough idea is that the equivariant signatures $\sigma_K(e^{2i(\gamma \pm \varepsilon)})$ are signed counts of points in the spaces of irreducible ρ with $\text{tr}(\rho(\mu)) = 2 \cos(\gamma \pm \varepsilon)$, and if $(\gamma, 0)$ is not a limit point then these spaces are cobordant as in [HK98], hence $\sigma_K(e^{i(\theta-2\varepsilon)}) = \sigma_K(e^{i(\theta+2\varepsilon)})$. We briefly indicate the details below, referring to [HK98] for definitions.

Letting $\sigma \in B_n$ be a braid with closure K and taking $\varepsilon > 0$ sufficiently small, so that in particular γ is the only root of $\Delta_K(e^{2it})$ in the interval $\gamma - \varepsilon \leq t \leq \gamma + \varepsilon$, the intersection

$$\hat{H} := \left(\hat{\Lambda}_n \cap \hat{\Gamma}_\sigma \right) \cap \bigcup_{\alpha \in [\gamma - \varepsilon, \gamma + \varepsilon]} \hat{H}_n^\alpha$$

in \hat{H}_n (as defined in [HK98]) is identified with the space of conjugacy classes of irreducible representations $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ satisfying

$$2 \cos(\gamma - \varepsilon) \leq \operatorname{tr}(\rho(\mu)) \leq 2 \cos(\gamma + \varepsilon).$$

The set \hat{H} is not quite compact because it contains a sequence which limits to the conjugacy class $[\rho_\gamma] \notin \hat{H}$. Supposing that $(\gamma, 0)$ is an isolated point, however, its preimage in \hat{H} is a union of connected components of \hat{H} and is contained in \hat{H}_n^γ , so we let $\hat{H}' \subset \hat{H}_n$ be the complement of these components. Then \hat{H}' is a compact cobordism from

$$\hat{\Lambda}_n^{\gamma-\varepsilon} \cap \tilde{\Gamma}_\sigma^{\gamma-\varepsilon} \subset \hat{H}_n^{\gamma-\varepsilon} \quad \text{to} \quad \hat{\Lambda}_n^{\gamma+\varepsilon} \cap \tilde{\Gamma}_\sigma^{\gamma+\varepsilon} \subset \hat{H}_n^{\gamma+\varepsilon},$$

which implies as in [HK98] that $h^{\gamma-\varepsilon}(K) = h^{\gamma+\varepsilon}(K)$. Now [HK98, Theorem 1.2] gives us $\sigma_K(e^{2i(\gamma-\varepsilon)}) = \sigma_K(e^{2i(\gamma+\varepsilon)})$, where σ_K denotes equivariant signature, and since $\Delta_K(t)$ has a root of odd order at $e^{2i\gamma}$, we have a contradiction.

Finally, we suppose that K is $SU(2)$ -averse with limit slope $r(K) = \frac{p}{q}$. As shown above, we can find a sequence of distinct points in the pillowcase image of $X^*(K)$ which limit to $(\gamma, 0)$, and by passing to a subsequence we can take them all to lie in the same path component. This path component must then be a nontrivial segment of the line of slope $-\frac{p}{q}$ passing through $(\gamma, 0)$, namely

$$p\alpha + q\beta \equiv p\gamma \pmod{2\pi},$$

but $p\gamma$ is not a rational multiple of 2π since $z = e^{2i\gamma}$ is not a root of unity. This contradicts Theorem 6.4, so K cannot be $SU(2)$ -averse after all. \square

Corollary 6.6. *If K has Alexander polynomial $\Delta_K(t) = at - (2a - 1) + at^{-1}$ for some integer $a \geq 2$, then K is not $SU(2)$ -averse.*

Proof. We have $\Delta_K(-1) < 0$, and $\Delta_K(t)$ is irreducible and not cyclotomic. \square

Example 6.7. The pretzel knot $K = P(-2, 3, 7)$ has Alexander polynomial

$$\Delta_K(t) = t^5 - t^4 + t^2 - t + 1 - t^{-1} + t^{-2} - t^{-4} + t^{-5},$$

which has a root of odd order on the unit circle since $\Delta_K(i) = -3$. Then $\Delta_K(t)$ is irreducible and not cyclotomic (in fact, $-t^5\Delta_K(-t)$ is the minimal polynomial of the smallest known Salem number: see [Smy15, Section 2]), so this root is not a root of unity. Proposition 6.5 thus ensures that $P(-2, 3, 7)$ is not $SU(2)$ -averse.

7. SMALL KNOTS

We recall that a knot $K \subset S^3$ is *small* if its complement does not contain any closed incompressible surfaces other than boundary-parallel tori. In this section we use the results of Section 6, together with work of Boyer and Zhang [BZ98], to develop further restrictions on the limit slopes of small $SU(2)$ -averse knots.

Theorem 7.1. *If K is a small, $SU(2)$ -averse knot, then its limit slope $r(K)$ is an integer. In this case, we have $\rho(\mu^{r(K)}\lambda) = \pm I$ for all but finitely many conjugacy classes of irreducible representations $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$.*

Proof. We write $r(K) = \frac{p}{q}$ with p and q relatively prime. Since K is small, every irreducible component of the $SL_2(\mathbb{C})$ character variety $X_{SL_2}(K)$ is a curve [CCG⁺94, Proposition 2.4]. One of these components, say X_0 , contains infinitely many distinct points which are the

images of $SU(2)$ -representations on a line of the form $p\alpha + q\beta = c$ but with distinct β -coordinates in the pillowcase, so $\text{tr}(\rho(\mu^p\lambda^q)) = 2\cos(c)$ at infinitely many points on the curve X_0 , hence $\text{tr}(\rho(\mu^p\lambda^q))$ is constant on X_0 . The image of X_0 in the $PSL_2(\mathbb{C})$ character variety of K is then a $\frac{p}{q}$ -curve in the terminology of Boyer-Zhang [BZ98] – we note that the traces of other slopes cannot also be constant on X_0 , since for example $\text{tr}(\rho(\lambda))$ is not constant – and the meridian of K is not a boundary slope [CGLS87, Theorem 2.0.3], so by [BZ98, Corollary 6.7], the slope $\frac{p}{q} = r(K)$ must be integral.


Letting $r = r(K)$, any nontrivial segment of a line $\beta = -r\alpha + c$ in the pillowcase image of $X^*(K)$ produces an r -curve X_0 as above, and clearly X_0 contains the character of an irreducible representation. Since the exterior of K is small and admits a cyclic filling of slope $\infty \neq r$, we apply [BZ98, Proposition 5.7(2)] to see that $\text{tr}(\rho(\mu^r\lambda)) = \pm 2$ on all of X_0 , hence the $SU(2)$ -representations which contribute to X_0 must satisfy $\rho(\mu^r\lambda) = \pm I$. Thus $\rho(\mu^r\lambda) = \pm I$ for every ρ with pillowcase image on the line $\beta = -r\alpha + c$.

All but finitely many isolated points in the pillowcase image of $X^*(K)$ belong to a nontrivial line segment, so it remains to be seen that an isolated point in the pillowcase can only be the image of finitely many conjugacy classes of $SU(2)$ representations. Suppose that infinitely many conjugacy classes $[\rho_n] \in R^*(K)$ have the same image (α, β) . The ρ_n remain pairwise non-conjugate as $SL_2(\mathbb{C})$ representations by [Kla91, Proof of Proposition 15], so they produce infinitely many points in $X_{SL_2}(K)$ and hence in some irreducible component X_1 of the $PSL_2(\mathbb{C})$ character variety. By [BZ98, Proposition 5.7], the function $f_\gamma = \text{tr}(\rho(\gamma))^2 - 4 : X_1 \rightarrow \mathbb{C}$ cannot be constant for two different slopes γ , but it is clearly constant for the slopes μ and λ and this is a contradiction. \square

The claim in Theorem 7.1 that $\rho(\mu^{r(K)}\lambda) = \pm I$ for representations along a nontrivial segment of $\beta = -r\alpha + c$ can also be proved using the A -polynomial as follows. Proposition 6.2 says that $M^rL - e^{ic}$ divides $A_K(M, L)$, so e^{ic} is a root of $A_K(1, L)$. Since K is small, the latter is equal to $\pm(L+1)^\alpha(L-1)^\beta L^\gamma$ for some integers α, β, γ by [CL96, Corollary 4.5], so its nonzero root e^{ic} must be ± 1 .

Example 7.2. Let K_n denote the twist knot shown in Figure 4, where $n \in \frac{1}{2}\mathbb{Z}$. Then K_n is a two-bridge knot, hence small, and moreover K_n is hyperbolic for all n other than $-1, -\frac{1}{2}, 0$, and $\frac{1}{2}$, for which it is the right-handed trefoil, unknot, unknot, and left-handed trefoil, respectively. We claim that no hyperbolic twist knot is $SU(2)$ -averse; since K_n is the mirror of $K_{-\frac{1}{2}-n}$, it suffices to consider $n \in \mathbb{Z}$.

Indeed, suppose for some integer $n \notin \{-1, 0\}$ that K_n is $SU(2)$ -averse with limit slope $r = r(K_n)$. Then just as in the proof of Theorem 7.1 we see that the $SL_2(\mathbb{C})$ character variety $X_{SL_2}(K_n)$ has an irreducible component X_0 , containing irreducible characters, for which the function $\text{tr}(\rho(\mu^r\lambda))|_{X_0}$ is constant. Burde [Bur90, Section 3] showed that all irreducible $SL_2(\mathbb{C})$ characters of $\pi_1(S^3 \setminus K_n)$ lie on a single irreducible curve in $X_{SL_2}(K)$, so this curve must be X_0 ; then since K_n is hyperbolic we see that X_0 must also contain the character of a discrete faithful $SL_2(\mathbb{C})$ representation, which exists by a theorem of Thurston [CS83, Proposition 3.1.1]. But then [CS84, Proposition 2] says that if γ is any nontrivial peripheral curve then the function $\text{tr}(\rho(\gamma))$ cannot be constant on X_0 , so we have a contradiction.

FIGURE 4. The twist knot K_n .

Theorem 7.3. *Let K be a small, $SU(2)$ -averse knot with boundary slope $r(K)$ and with Alexander polynomial*

$$\Delta_K(t) = \sum_{j=-d}^d a_j t^j,$$

normalized so that $\Delta_K(t) = \Delta_K(t^{-1})$ and $\Delta_K(1) = 1$. Then

$$\det(K) \leq \sum_{j=-d}^d |a_j| \leq \text{rank}(KHI(K)) \leq |r(K)| - 1,$$

where $KHI(K)$ is the instanton knot homology of K .

Proof. The leftmost two inequalities come from the fact that $KHI(K)$ categorifies the Alexander polynomial [KM10a, Lim10], meaning that it has a decomposition

$$KHI(K) = \bigoplus_{j \in \mathbb{Z}} KHI(K, j)$$

into canonically $\mathbb{Z}/2\mathbb{Z}$ -graded summands satisfying

$$\Delta_K(t) = - \sum_{j \in \mathbb{Z}} \chi(KHI(K, j)) t^j,$$

and so $\det(K) = |\Delta_K(-1)| \leq \sum_j |a_j| \leq \sum_j \text{rank}(KHI(K, j)) = \text{rank}(KHI(K))$. Thus we only need to prove that $\text{rank}(KHI(K)) \leq |r(K)| - 1$.

This last inequality will be proved by an application of Theorem 4.8. It suffices to find an area-preserving isotopy of the pillowcase which fixes the four corners and carries $C = \{\alpha = \frac{\pi}{2}\}$ to a curve C' which avoids both the image of $X^*(K)$ and the points $(\alpha, 0)$ where $\Delta_K(e^{2i\alpha}) = 0$. If C' intersects the line $\beta \equiv 0 \pmod{2\pi}$ transversely in at most $|r(K)| - 1$ points, then this will prove the desired inequality.

Since K is small, all but finitely many conjugacy classes of irreducible representations $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ satisfy $\rho(\mu^r \lambda) = \pm I$ where $r = r(K)$, and hence their images (α, β) in the pillowcase satisfy $r\alpha + \beta \equiv 0 \pmod{\pi}$. We take $\varepsilon > 0$ so that the irreducible characters all satisfy $\varepsilon < \alpha < \pi - \varepsilon$. We then let U be a δ -neighborhood of the set

$$\left\{ (\alpha, \beta) \mid r\alpha + \beta \in \pi\mathbb{Z}, \frac{\varepsilon}{2} \leq \alpha \leq \frac{\pi}{2} \right\} \cup \left\{ \left(\frac{\pi}{2}, \beta \right) \mid \beta \in \mathbb{R}/2\pi\mathbb{Z} \right\},$$

where $0 < \delta < \frac{\varepsilon}{2}$, and let Z_0 denote the component of ∂U contained in the region $\alpha < \frac{\pi}{2}$. The curves corresponding to different choices of δ are all disjoint, so only finitely many of them can pass through the images of irreducible characters or through points $(\alpha, 0)$ where $\Delta_K(e^{2i\alpha}) = 0$. Taking δ sufficiently small thus ensures that Z_0 avoids all such points.

Next, we perturb Z_0 to a smooth curve Z which separates $X(T^2)$ into two components of equal area, by rounding corners and then pushing some segment of Z_0 slightly into the $\alpha > \frac{\pi}{2}$ region of the pillowcase away from the lines $r\alpha + \beta \equiv 0 \pmod{\pi}$ and $\beta \equiv 0$




FIGURE 5. A perturbation Z of the line $C = \{\alpha = \frac{\pi}{2}\}$ in the pillowcase, shown for $r = 4$ (left) and $r = 5$ (right). The diagonal dotted lines are the points where $r\alpha + \beta \equiv 0 \pmod{\pi}$.

(mod 2π), as shown in Figure 5. This is possible because the difference in area between the two components of $X(T^2) \setminus Z_0$ approaches zero as δ does, and once again it can easily be arranged to avoid all images of irreducible characters. We also arrange that Z crosses the line $\beta \equiv 0 \pmod{2\pi}$ transversely. It does so twice near each point $(\alpha, 0)$ on the pillowcase with $r\alpha \in \pi\mathbb{Z}$ and $0 < \alpha < \frac{\pi}{2}$, and once near $(\frac{\pi}{2}, 0)$, for a total of $|r| - 1$ points if r is even and $|r|$ points if r is odd.

In the case where r is odd, we can perturb Z further to remove two of its points of intersection with $\beta \equiv 0 \pmod{2\pi}$, as shown in Figure 6. The idea is that in a fundamental domain $[0, \pi] \times [0, 2\pi]$ for the pillowcase, if we assume for convenience that $r > 0$ (the case $r < 0$ is nearly identical), then there is a triangle with vertices

$$\left(\frac{r-1}{2r}\pi, 2\pi\right), \quad \left(\frac{\pi}{2}, 2\pi\right), \quad \left(\frac{\pi}{2}, \frac{3\pi}{2}\right),$$

whose area is $\frac{\pi^2}{8r}$, and an arc of Z which intersects this triangle in two points along the line $\beta = 2\pi$. We can push this arc up across the line $\beta = 2\pi$ to remove these two points, and the total area lost on that side of Z (i.e., the side incident to $\alpha = 0$) is less than $\frac{\pi^2}{8r}$. We then offset this change in area by simultaneously pushing Z further into the interior of the quadrilateral bounded by the points

$$\left(\frac{r+1}{2r}\pi, 0\right), \quad \left(\frac{\pi}{2}, \frac{\pi}{2}\right), \quad \left(\frac{\pi}{2}, \frac{3\pi}{2}\right), \quad \left(\frac{r+3}{2r}\pi, 0\right),$$

whose area is $\frac{\pi^2}{r}$, so that Z still divides $X(T^2)$ into two halves of equal area.

The end result is that the curve $C' = Z$ intersects $\beta \equiv 0 \pmod{2\pi}$ in $|r| - 1$ points if r is even and $|r| - 2$ points if r is odd, and it is easy to construct an area-preserving isotopy of $X(T^2)$ which fixes the four corners and carries it to C , so we conclude that $\text{rank}(KHI(K)) \leq |r| - 1$. \square

Although Proposition 6.5 does not say anything useful about knots whose Alexander polynomials have zeros at roots of unity, we can use such zeros to deduce useful information for small knots as follows.




FIGURE 6. If r is odd, we can perturb the curve Z from Figure 5 so that it only intersects the curve $\beta \equiv 0 \pmod{2\pi}$ in $|r| - 2$ points.

Proposition 7.4. *Let K be a small knot which is $SU(2)$ -averse. If the Alexander polynomial $\Delta_K(t)$ has a root of odd order at $t = e^{i \cdot 2\pi p/q}$, where p and q are relatively prime integers, then the limit slope $r(K)$ is a multiple of q .*

Proof. Let $r = r(K)$, and note that the line $r\alpha + \beta \equiv r \cdot \frac{\pi p}{q} \pmod{2\pi}$ has slope $-r$ and passes through the image $(\frac{\pi p}{q}, 0)$ of the character of the reducible representation $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ with $\rho(\mu) = \text{diag}(e^{i\pi p/q}, e^{-i\pi p/q})$. Arguing exactly as in the proof of Proposition 6.5, we see that the pillowcase image of $X^*(K)$ contains a nontrivial segment of this line. By Theorem 7.1, we have $r \cdot \frac{\pi p}{q} \equiv 0 \pmod{\pi}$, and so q divides r as claimed. \square

Proposition 7.5. *If K is a small, $SU(2)$ -averse knot, then $|r(K)| \geq 6$.*

Proof. Suppose that $|r(K)| \leq 5$. Then Theorem 7.3 says that $\text{rank}(KHI(K)) \leq 4$, and in fact the rank of $KHI(K)$ is odd (since it categorifies the Alexander polynomial and $\Delta_K(1) = 1$ is odd), so we have $\text{rank}(KHI(K)) \leq 3$. By assumption K is not the unknot, so we must have $\text{rank}(KHI(K)) = 3$.

Kronheimer and Mrowka proved that $\text{rank } KHI(K, g) \geq 1$ [KM10b, Proposition 7.16], where $g \geq 1$ is the Seifert genus of K . Since $KHI(K, j) \cong KHI(K, -j)$ for all j , it follows that $KHI(K, j)$ has rank one for each of $j = -g, 0, g$ and rank zero for all other j , and this immediately implies that

$$\Delta_K(t) = t^g - 1 + t^{-g}.$$

But then $\Delta_K(t)$ has a simple root at $t = e^{i \cdot 2\pi/(6g)}$, so Proposition 7.4 tells us that $|r(K)|$, which is an integer between 1 and 5, is a multiple of $6g \geq 6$, which is absurd. \square

We remark that Proposition 7.5 is sharp, since the right- and left-handed trefoils are $SU(2)$ -averse with limit slope ± 6 .

8. $SU(2)$ -CYCLIC SURGERIES FROM THE PILLOWCASE

In this section we prove a converse to Theorem 3.5. Our goal is to say that if the pillowcase image of $X^*(K)$, the irreducible part of the character variety, is contained in finitely many lines of rational slope $-r$, then K must be $SU(2)$ -averse with limit slope r . We will use the following lemma to prove such a result.

Lemma 8.1. *Let $r, c \in \mathbb{R}$ be constants, and fix a rational number $\frac{p}{q} \neq r$ with $q > 0$ and p relatively prime to q . Let $\{x\} = x - \lfloor x \rfloor$ denote the fractional part of x . If*

$$0 \leq \frac{q}{2} \left(\frac{p}{q} - r \right) + \left\{ \frac{qc}{2\pi} \right\} \leq 1,$$

then no point on the line $\beta = -r\alpha + c$, $0 < \alpha < \pi$, satisfies $p\alpha + q\beta \equiv 0 \pmod{2\pi}$.

Proof. Let $k = \lfloor \frac{qc}{2\pi} \rfloor$. Then the given inequalities are equivalent to

$$k \leq \frac{q}{2} \left(\frac{p}{q} - r \right) + \frac{qc}{2\pi} \leq k + 1,$$

which can be rearranged to get

$$\frac{-p\pi + 2\pi k}{q} \leq -r\pi + c \leq \frac{-p\pi + 2\pi(k+1)}{q}.$$

Thus we have a pair of linear inequalities in α of the form

$$\frac{-p\alpha + 2\pi k}{q} \leq -r\alpha + c \leq \frac{-p\alpha + 2\pi(k+1)}{q},$$

which are satisfied at $\alpha = 0$ (since our choice of k implies that $\frac{2\pi k}{q} \leq c < \frac{2\pi(k+1)}{q}$) and at $\alpha = \pi$, and hence for all α in between. In fact, each of these inequalities must be strict for $0 < \alpha < \pi$: otherwise, since $\frac{p}{q} \neq r$, the corresponding inequality would be violated on one side of the α in question. Thus $\beta = -r\alpha + c$ lies strictly between $\beta = \frac{-p\alpha + 2\pi k}{q}$ and $\beta = \frac{-p\alpha + 2\pi(k+1)}{q}$ for $0 < \alpha < \pi$, and so it cannot contain any point on the lines $p\alpha + q\beta \equiv 0 \pmod{2\pi}$, as claimed. \square

Theorem 8.2. *Let K be a nontrivial knot, and suppose that there is a constant $r \in \mathbb{Q}$ such that the pillowcase image of $X^*(K)$ is contained in the union of finitely many isolated points and finitely many nontrivial segments of lines $\beta = -r\alpha + c_i$. Then there is some $N \geq 1$ such that $(r + \frac{1}{kN})$ -surgery on K is SU(2)-cyclic for all but finitely many $k \in \mathbb{Z}$. In particular, K is SU(2)-averse.*

The hypotheses of Theorem 8.2 are satisfied by all SU(2)-averse knots: this is guaranteed for some $r \in \mathbb{RP}^1$ by Theorem 3.5, and then Theorems 4.2 and 5.4 assert that r is rational.

Proof. According to Theorem 6.3, the constants c_i are all rational multiples of 2π . Let $n \geq 1$ be an integer such that nr and each of the $\frac{nc_i}{\pi}$ are integers, and let $m = nr$ and $N = n^2$. For a given slope $\frac{p}{q} \in \mathbb{Q}$ not equal to r , with $q > 0$ and p relatively prime to q , if the inequalities

$$0 \leq \frac{q}{2} \left(\frac{p}{q} - r \right) + \left\{ \frac{qc_i}{2\pi} \right\} \leq 1$$

are satisfied for all i , then Lemma 8.1 says that no point (α, β) on one of the lines $\beta = -r\alpha + c_i$ which is the image of an irreducible $\rho : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ can satisfy $p\alpha + q\beta \equiv 0 \pmod{2\pi}$ (recall that $0 < \alpha < \pi$, by Proposition 2.1). We will take $(p, q) = (knm + 1, kn^2)$ for any integer $k \geq 1$, noting that p and q are relatively prime since $km^2q - (knm - 1)p = 1$.

We compute for each i that

$$\begin{aligned} \frac{q}{2} \left(\frac{p}{q} - r \right) + \left\{ \frac{qc_i}{2\pi} \right\} &= \frac{kn^2}{2} \left(\frac{knm+1}{kn^2} - \frac{m}{n} \right) + \left\{ kn \cdot \frac{nc_i}{2\pi} \right\} \\ &= \frac{1}{2} + \left\{ \frac{1}{2} \cdot kn \cdot \frac{nc_i}{\pi} \right\}, \end{aligned}$$

which is either $\frac{1}{2}$ or 1, so the irreducible representations on the lines $\beta = -r\alpha + c_i$ do not produce irreducible representations of the $\frac{kmn+1}{kn^2}$ -surgery on K .

We are left with finitely many isolated points (α_j, β_j) ; letting $c'_j = \beta_j + r\alpha_j$, we may assume without loss of generality that $c'_j \notin 2\pi\mathbb{Q}$, since otherwise we could have just added the line $\beta = -r\alpha + c'_j$ to the collection of lines considered above, possibly at the cost of taking a larger value of n . If for some $k \geq 1$ we have $(kmn+1)\alpha_j + kn^2\beta_j \in 2\pi\mathbb{Z}$, then this together with $kn^2(\frac{m}{n}\alpha_j + \beta_j - c'_j) = 0$ implies that $\alpha_j + kn^2c'_j \in 2\pi\mathbb{Z}$. For fixed j , there is at most one k for which this can be satisfied since c'_j is not a rational multiple of 2π , and hence each (α_j, β_j) produces an irreducible representation of at most one $\frac{kmn+1}{kn^2}$ -surgery on K . Therefore the surgery on K of slope $\frac{kmn+1}{kn^2} = r + \frac{1}{kN}$ is $SU(2)$ -cyclic for all but finitely many $k \geq 1$.

Next, the mirror \overline{K} satisfies the same hypotheses with $-r$ and $2\pi - c_i$ in place of r and c_i by Proposition 2.4, so we can take the same value of n (and hence N) as above and conclude that all but finitely many $(-r + \frac{1}{lN})$ -surgeries on \overline{K} are $SU(2)$ -cyclic when $l \geq 1$. These are the $(r - \frac{1}{lN})$ -surgeries on K up to reversing orientation, so taking $k = -l$ we see that all but finitely many $(r + \frac{1}{kN})$ -surgeries on K with $k \leq -1$ are $SU(2)$ -cyclic as well. \square

If K is small and $SU(2)$ -averse, then by Theorem 7.1 we have $r(K) \in \mathbb{Z}$ and $c_i \in \pi\mathbb{Z}$, so if there are no isolated points in the pillowcase image of $X^*(K)$ then we can take $n = 1$ and hence $N = 1$ in the proof of Theorem 8.2. In this case, we conclude that every slope $r(K) + \frac{1}{k}$ (with $k \in \mathbb{Z}$ nonzero) is an $SU(2)$ -cyclic slope for K .

Corollary 8.3. *If K_1, K_2, \dots, K_m are $SU(2)$ -averse knots which all have the same limit slope, then they have infinitely many $SU(2)$ -cyclic surgery slopes in common.*

Proof. Let $r \in \mathbb{Q}$ be the common limit slope and take $N_1, N_2, \dots, N_m \geq 1$ such that for each i , all but finitely many of the $(r + \frac{1}{kN_i})$ -surgeries on K_i are $SU(2)$ -cyclic. Then for all but finitely many k , all of the $(r + \frac{1}{kN})$ -surgeries on the various K_i are $SU(2)$ -cyclic, where $N = \text{lcm}(N_1, N_2, \dots, N_m)$. \square

9. RELATION TO INSTANTON L-SPACES

Let Y be a closed, connected, oriented 3-manifold, and let $I^\#(Y)$ be the singular instanton knot homology $I^\#(Y, \emptyset) = I^\natural(Y, U)$ of the empty link in Y , as defined by Kronheimer and Mrowka [KM11]. This is a $\mathbb{Z}/4\mathbb{Z}$ -graded abelian group, and if Y is a rational homology sphere then it has Euler characteristic $|H_1(Y; \mathbb{Z})|$, as shown by Scaduto [Sca15, Corollary 1.4]. Thus we have a rank inequality

$$\text{rank}(I^\#(Y)) \geq |H_1(Y; \mathbb{Z})|,$$

and if equality holds then we say that Y is an *instanton L-space*.

Theorem 9.1. *If K is $SU(2)$ -averse with limit slope $r = r(K)$, then $S_s^3(K)$ is an instanton L-space for all rational*

$$s \in \begin{cases} [[r] - 1, \infty), & r > 0 \\ (-\infty, [r] + 1], & r < 0. \end{cases}$$

In either case, $S_r^3(K)$ is an instanton L-space.

Remark 9.2. Conjecturally, being an instanton L-space is the same as being an L-space in Heegaard Floer homology. This would imply for $SU(2)$ -averse knots K that all of the nonzero coefficients of $\Delta_K(t)$ are ± 1 [OS05], that K is fibered [Ni07], and that either K or its mirror is strongly quasipositive [Hed10]. The proof of [BS16, Theorem 1.10] shows that negative surgeries on strongly quasipositive knots are not instanton L-spaces, so Theorem 9.1 implies that a strongly quasipositive knot which is $SU(2)$ -averse has positive limit slope.

Proof of Theorem 9.1. Assume for now that $r(K) > 0$, and write $r(K) = \frac{p}{q}$ for coprime positive integers p and q . Theorem 8.2 provides an integer $N > 0$ such that for all but finitely many $k > 0$, $r(K) - \frac{1}{kN}$ is an $SU(2)$ -cyclic surgery slope for K . (It is also at least $r(K) - 1 > 1$, hence positive.) In particular, this is true for all but finitely many multiples $k = qi$ of q , in which case the slopes in question are

$$\frac{p}{q} - \frac{1}{qi \cdot N} = \frac{pNi - 1}{qNi}$$

for some $i > 0$. The numerator $pNi - 1$ is prime for infinitely many values of i by Dirichlet's theorem on primes in arithmetic progressions, so K must have a positive $SU(2)$ -cyclic surgery slope of the form $\frac{a}{b} < \frac{p}{q}$, where a is prime.

If $S_{a/b}^3(K)$ is not an instanton L-space, then [BS16, Corollary 4.8] says that either $S_{a/b}^3(K)$ is not $SU(2)$ -cyclic or some zero ζ of $\Delta_K(t^2)$ is an a th root of unity. The former is false by construction, and the latter cannot hold either: the minimal polynomial $\Phi_a(t) = 1 + t + t^2 + \dots + t^{a-1}$ of ζ also divides $\Delta_K(t^2)$, and so $\Phi_a(1) = a$ must divide $\Delta_K(1^2) = 1$, which is impossible. Thus $S_{a/b}^3(K)$ is an instanton L-space.

Now if $n = \lfloor \frac{a}{b} \rfloor$, then $S_s^3(K)$ is an instanton L-space for all $s \geq n$ by [BS16, Theorem 4.20]. Since n is an integer which is strictly less than $r(K)$, we have $n \leq [r(K)] - 1$, and this completes the proof in the case $r(K) > 0$.

If instead K has negative limit slope, then the mirror \bar{K} satisfies $r(\bar{K}) = -r(K) > 0$. The above argument shows that $S_s^3(\bar{K}) = -S_{-s}^3(K)$ is an instanton L-space whenever

$$s \geq [-r(K)] - 1 = -[r(K)] - 1,$$

or equivalently whenever $-s \leq [r(K)] + 1$. Since the property of being an instanton L-space is preserved under orientation reversal, this proves the case $r(K) < 0$. \square

It is generally hard to show that a knot does not have any instanton L-space surgeries, but when we can do so it follows by Theorem 9.1 that the knot in question is not $SU(2)$ -averse.

Theorem 9.3. *If a nontrivial knot $K \subset S^3$ is smoothly slice, then it has no nontrivial instanton L-space surgeries. In particular, K is not $SU(2)$ -averse.*

Proof. Suppose that K is slice and that $S_r^3(K)$ is an instanton L-space for some rational r ; by taking mirror images as needed we can assume that $r > 0$. Then [BS16, Theorem 4.20] says that $S_n^3(K)$ is an instanton L-space for all integers $n \geq r$.

Letting $m \geq 1$ be an integer, we now apply Floer's surgery exact triangle

$$\dots \rightarrow I^\#(S^3) \rightarrow I^\#(S_m^3(K)) \rightarrow I^\#(S_{m+1}^3(K)) \rightarrow \dots$$

as stated in [Sca15, Section 7.5], cf. also [Flo90, BD95]. (This sequence involves twisted coefficients, but following the third row of [Sca15, Figure 1] we can place the nontrivial coefficients in the first term; the twisted group $I^\#(S^3; \mu)$ is isomorphic to $I^\#(S^3)$, since $[\mu] = 0$ in $H_1(S^3; \mathbb{Z}/2\mathbb{Z})$.) Each of these maps is induced by a 2-handle cobordism, and the cobordism from S^3 to $S_m^3(K)$ contains a smoothly embedded 2-sphere S of self-intersection $m \geq 1$, obtained by gluing a slice disk for K to the core of the 2-handle.

Since the sphere S has positive self-intersection, the map $I^\#(S^3) \rightarrow I^\#(S_m^3(K))$ necessarily vanishes. This can be proved directly from the results of [MMR94], or indirectly from [MMR94, Corollary 16.0.2] by using S to construct a torus of self-intersection at least 2: first, if $m = 1$ then we patch two copies of S together to get a sphere S' of self-intersection 4, and then we attach a handle to either S or S' to get the desired torus. Thus the above triangle splits, and since $I^\#(S^3) \cong \mathbb{Z}$ we have

$$\text{rank}(I^\#(S_{m+1}^3(K))) = \text{rank}(I^\#(S_m^3(K))) + 1,$$

hence $\text{rank}(I^\#(S_m^3(K))) = m + (\text{rank}(I^\#(S_1^3(K))) - 1)$ for all $m \geq 1$ by induction.

We conclude that $S_n^3(K)$ is an instanton L-space for each integer $n \geq r$ if and only if $S_1^3(K)$ is. By [BS16, Theorem 4.20], the latter can only be true if K has Alexander polynomial $t - 1 + t^{-1}$ or $-t + 3 - t^{-1}$. But then K violates the Fox-Milnor condition, which asserts that slice knots satisfy $\Delta_K(t) = f(t)f(t^{-1})$ for some polynomial $f(t)$ with integer coefficients [FM66], so it cannot be slice and we have a contradiction. \square

Proposition 9.4. *If K is $SU(2)$ -averse with smooth slice genus 1, then either $S_1^3(K)$ or $S_{-1}^3(K)$ is an instanton L-space depending on whether $r(K)$ is positive or negative. Thus K has Seifert genus 1 and its Alexander polynomial is either $t - 1 + t^{-1}$ or $-t + 3 - t^{-1}$.*

Proof. We arrange for $r(K) > 0$ by replacing K with its mirror as necessary, and then repeat the proof of Theorem 9.3: each $S_n^3(K)$ is an instanton L-space for integers $n \geq r(K)$, and so by the surgery exact triangle and induction, $S_1^3(K)$ must be as well. The 2-handle cobordisms from S^3 to $S_m^3(K)$ now have an embedded torus (rather than sphere) of self-intersection $m \geq 1$, and for $m \geq 2$ we can still apply [MMR94, Corollary 16.0.2] to see that the cobordism maps on $I^\#$ vanish.

The only change needed is in the case $m = 1$, where we prove the vanishing of the cobordism map by using the adjunction inequality of [KM95] instead. This requires some attention, since [KM95] assumes the ambient manifold is closed with $b^+ \geq 3$. However, the same arguments as in [KM95] prove the vanishing of maps associated to cobordisms which contain smooth surfaces of odd genus g and self-intersection $n \not\equiv 0 \pmod{4}$ with $\frac{n}{2} + 2 < 2g - 2$ (see [KM95, Section 6(iii)]). We take five copies in general position of a torus T with self-intersection 1 and replace a neighborhood of each of the $\binom{5}{2} = 10$ transverse intersection points with an annulus to get an embedded surface Σ in the cobordism with $n = (5T) \cdot (5T) = 25$ and with $\chi(\Sigma) = 5\chi(T^2) - 10(2) = -20$, hence genus $g = 11$, and so the cobordism map on $I^\#$ is zero.

Thus if $r(K) > 0$ then $S_1^3(K)$ is an instanton L-space, and if $r(K) < 0$ then the same is true of $S_1^3(\bar{K}) = -S_{-1}^3(K)$, hence also of $S_{-1}^3(K)$. We now apply [BS16, Theorem 4.20] to K or \bar{K} to draw the desired conclusions about its genus and Alexander polynomial. \square

We expect that the right-handed trefoil is the only nontrivial knot $K \subset S^3$ for which $S^3_1(K)$ is an instanton L-space. (Indeed, Proposition 9.4 proves that 1-surgery on this trefoil is an instanton L-space.) If this were the case, then Proposition 9.4 would show that the trefoils are the only $SU(2)$ -averse knots with slice genus at most 1.

Example 9.5. We consider the twist knots K_n of Example 7.2, assuming $n \in \mathbb{Z}$ without loss of generality. These all have Seifert genus and hence smooth slice genus at most 1, and Alexander polynomial $\Delta_{K_n}(t) = -nt + (2n + 1) - nt^{-1}$, so by Proposition 9.4 these cannot possibly be $SU(2)$ -averse except when $n = -1, 0, 1$. The case $n = 1$ is the figure eight, which cannot be $SU(2)$ -averse because it is amphichiral: otherwise we would have $r(K_1) = r(\overline{K_1}) = -r(K_1)$ and hence $r(K_1) = 0$, which is impossible. Thus we see once again that no hyperbolic twist knot is $SU(2)$ -averse.

10. SATELLITES AND CABLING

Let $P \subset S^1 \times D^2$ be an oriented knot in the solid torus which does not lie in any embedded 3-ball, and let $K \subset S^3$ be a knot with tubular neighborhood $N(K)$ and peripheral curves $\mu, \lambda \in \partial N(K)$. Then we can take a homeomorphism

$$\varphi : S^1 \times D^2 \xrightarrow{\sim} N(K) \subset S^3$$

which sends $\{\text{pt}\} \times \partial D^2$ to μ and $S^1 \times \{\text{pt}\} \subset S^1 \times \partial D^2$ to λ , and we define the *satellite* $P(K) \subset S^3$ to be the image $\varphi(P)$. The knots P and K are called the *pattern* and the *companion*, respectively, and P is *nontrivial* if it is not isotopic to the core of $S^1 \times D^2$. The *winding number* of P is the unique integer w such that P represents an element $w[S^1 \times \{\text{pt}\}]$ of $H^1(S^1 \times D^2)$. From now on U will denote the unknot.

The following is a result of Silver and Whitten, stated here in a slightly stronger form.

Proposition 10.1 ([SW06, Proposition 3.4]). *Given any pattern P and companion K , there is a surjection*

$$\psi : \pi_1(S^3 \setminus P(K)) \rightarrow \pi_1(S^3 \setminus P(U))$$

which preserves meridians and longitudes, i.e. such that $\psi(\mu_{P(K)}) = \mu_{P(U)}$ and $\psi(\lambda_{P(K)}) = \psi(\lambda_{P(U)})$.

Proof. We can write the knot group of $P(K)$ as an amalgamated free product

$$\pi_1(S^3 \setminus P(K)) = \pi_1(S^3 \setminus K) *_{\mathbb{Z}^2} \pi_1((S^1 \times D^2) \setminus P),$$

where generators of $\mathbb{Z}^2 = \pi_1(\partial N(K))$ identify a meridian and longitude of K with the curves $\{\text{pt}\} \times \partial D^2$ and $S^1 \times \{\text{pt}\}$ on $S^1 \times \partial D^2$ respectively. Silver and Whitten argue that taking the quotient of this product by the normal closure \mathcal{N} of the commutator subgroup of $\pi_1(S^3 \setminus K)$ reduces the first factor to \mathbb{Z} , generated by $\mu_K \sim (\{\text{pt}\} \times \partial D^2)$, and it sends the element $\lambda_K \sim (S^1 \times \{\text{pt}\})$ to the identity, so the quotient is the knot group of $P(U)$ and ψ is defined as the quotient map.

It is clear from the construction that $\psi(\mu_{P(K)})$ is still a meridian of $P(U)$. To see the analogous claim for $\lambda_{P(K)}$, we take an embedded surface Σ in $S^1 \times D^2$ with boundary $P \sqcup (S^1 \times \underline{w})$, where $\underline{w} \subset \partial D^2$ is a set of oriented points, and let $\lambda_P = \Sigma \cap \partial N(P) \subset (S^1 \times D^2) \setminus N(P)$. Then $\varphi(\lambda_P)$ is a longitude of any satellite $P(K)$, since the image $\varphi(S^1 \times \underline{w})$ bounds a collection of parallel Seifert surfaces for K . When we identify $\pi_1(S^3 \setminus P(U))$ as the quotient

$$\pi_1(S^3 \setminus P(K)) / \mathcal{N} \cong \pi_1((S^1 \times D^2) \setminus P) / \langle S^1 \times \{\text{pt}\} \rangle,$$

it follows that both $\lambda_{P(K)}$ and $\lambda_{P(U)}$ are represented by the same curve $\lambda_P \subset (S^1 \times D^2) \setminus N(P)$, and so $\psi(\lambda_{P(K)}) = \lambda_{P(U)}$. \square

The following are straightforward consequences of Proposition 10.1.

Proposition 10.2. *Let $P \subset S^1 \times D^2$ be a pattern and $K \subset S^3$ any knot.*

- (1) *The image of $X^*(P(U))$ in the pillowcase is a subset of the image of $X^*(P(K))$.*
- (2) *The $SU(2)$ -cyclic surgery slopes for $P(K)$ are all $SU(2)$ -cyclic slopes for $P(U)$.*
- (3) *If $P(K)$ is $SU(2)$ -averse and $P(U)$ is not the unknot, then $r(P(K)) = r(P(U))$.*

Proof. (1) If an irreducible $\rho : \pi_1(S^3 \setminus P(U)) \rightarrow SU(2)$ has pillowcase coordinates (α, β) , then so does $\tilde{\rho} = \rho \circ \psi : \pi_1(S^3 \setminus P(K)) \rightarrow SU(2)$, since $\psi(\mu_{P(K)}) = \mu_{P(U)}$ and $\psi(\lambda_{P(K)}) = \lambda_{P(U)}$. We have $\text{Im } \tilde{\rho} = \text{Im } \rho$ since ψ is surjective, so $\tilde{\rho}$ is irreducible as well.

(2) If $\frac{m}{n}$ -surgery on $P(K)$ is $SU(2)$ -cyclic then the line $m\alpha + n\beta \equiv 0 \pmod{2\pi}$ is disjoint from the image of $X^*(P(K))$, and hence also from the image of $X^*(P(U))$ by (1).

(3) The set of $SU(2)$ -cyclic slopes for $P(K)$ has limit point $r(P(K))$ by Theorem 3.2, hence so does the set of $SU(2)$ -cyclic slopes for $P(U)$ by (2). If $P(U)$ is not the unknot, then this limit point must therefore be $r(P(U))$, again by Theorem 3.2. \square

We can also prove more about satellites with nonzero winding number, as follows.

Proposition 10.3. *Let P be a satellite with winding number $w \neq 0$. If some r -surgery on $P(K)$ is $SU(2)$ -cyclic, then so is $\frac{r}{w^2}$ -surgery on K . In particular, if K is nontrivial and $P(K)$ is $SU(2)$ -averse, then K is also $SU(2)$ -averse, and $r(P(K)) = w^2 r(K)$.*

Proof. We will use the standard identification

$$S^3 \setminus P(K) = (S^3 \setminus N(K)) \cup ((S^1 \times D^2) \setminus P)$$

to construct representations $\rho \in R(P(K))$ whose restrictions to $S^3 \setminus N(K)$ are irreducible (so that ρ is also irreducible), but whose restrictions to $(S^1 \times D^2) \setminus P$ are abelian. Let μ_P, λ_P denote peripheral curves for P in $\partial N(P) \subset S^1 \times D^2$, so that μ_P bounds a meridian of P and the image $\varphi(\lambda_P) \subset \partial(S^3 \setminus N(P(K)))$ is a longitude of $P(K)$. We note that $[\lambda_P] = w[S^1 \times \{\text{pt}\}]$ as elements of $H_1((S^1 \times D^2) \setminus N(P))$.

Let $\rho_K : \pi_1(S^3 \setminus N(K)) \rightarrow SU(2)$ be an irreducible representation, where up to conjugacy we can write

$$\rho_K(\mu_K) = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{-i\alpha} \end{pmatrix}, \quad \rho_K(\lambda_K) = \begin{pmatrix} e^{i\beta} & 0 \\ 0 & e^{-i\beta} \end{pmatrix}$$

for some pair (α, β) . By the Mayer-Vietoris sequence we have $H^1((S^1 \times D^2) \setminus N(P)) \cong \mathbb{Z}^2$, with generators $[\mu_P]$ and $[S^1 \times \{\text{pt}\}]$ where $\text{pt} \in \partial D^2$. We know that $[\lambda_P] = w[S^1 \times \{\text{pt}\}]$, and similarly by examining a disk $\{\text{pt}\} \times D^2$ which intersects P transversely in w points (with sign) we see that $w[\mu_P] = [\{\text{pt}\} \times \partial D^2]$. So if we define an abelian representation

$$\rho_P : \pi_1((S^1 \times D^2) \setminus N(P)) \xrightarrow{\text{ab}} H_1((S^1 \times D^2) \setminus N(P)) \rightarrow SU(2)$$

by the formulas

$$\rho_P(\mu_P) = \begin{pmatrix} e^{i(\alpha+2\pi k)/w} & 0 \\ 0 & e^{-i(\alpha+2\pi k)/w} \end{pmatrix}, \quad \rho_P(S^1 \times \{\text{pt}\}) = \begin{pmatrix} e^{i\beta} & 0 \\ 0 & e^{-i\beta} \end{pmatrix}$$

for some integer k , then we have $\rho_P(\{\text{pt}\} \times \partial D^2) = \rho_K(\mu_K)$ and $\rho_P(S^1 \times \{\text{pt}\}) = \rho_K(\lambda_K)$. We glue ρ_K and ρ_P to get the desired representation $\rho \in R^*(P(K))$, with peripheral curves

$\mu = \varphi(\mu_P)$ and $\lambda = \varphi(\lambda_P)$ satisfying

$$\rho(\mu) = \begin{pmatrix} e^{i(\alpha+2\pi k)/w} & 0 \\ 0 & e^{-i(\alpha+2\pi k)/w} \end{pmatrix}, \quad \rho(\lambda) = \begin{pmatrix} e^{i\beta w} & 0 \\ 0 & e^{-i\beta w} \end{pmatrix},$$

and it follows that ρ has image $(\frac{\alpha+2\pi k}{w}, \beta w \pmod{2\pi})$ in the pillowcase.

Now suppose that some $\frac{p}{q}$ -surgery on $P(K)$ is $SU(2)$ -cyclic, where p and q are relatively prime. Let $d = \gcd(p, w^2)$, so that $\frac{p}{d}$ and $\frac{qw^2}{d}$ are also relatively prime. If the $\frac{p/d}{qw^2/d}$ -surgery on K is not $SU(2)$ -cyclic, then there is some irreducible $\rho_K \in R^*(K)$ with pillowcase image (α, β) satisfying

$$\frac{p}{d}\alpha + \frac{qw^2}{d}\beta = 2\pi n$$

for some integer n . The corresponding $\rho \in R^*(P(K))$ has image $(\frac{\alpha+2\pi k}{w}, \beta w)$, satisfying

$$p \left(\frac{\alpha + 2\pi k}{w} \right) + q(\beta w) = \frac{2\pi d}{w} (n + k(p/d)).$$

But $\frac{p}{d}$ is invertible modulo $\frac{w^2}{d}$ since these are coprime integers, so we can choose k so that $n + k(\frac{p}{d})$ is a multiple of $\frac{w^2}{d}$, and then the right side is a multiple of $2\pi w$. The representation ρ then descends to an irreducible representation of the $\frac{p}{q}$ -surgery on $P(K)$, which is impossible, so we conclude that $\frac{p/d}{qw^2/d} = \frac{p}{qw^2}$ is an $SU(2)$ -cyclic surgery slope for K after all. \square

10.1. Connected sums. We can now show that all $SU(2)$ -averse knots are prime.

Theorem 10.4. *If K_1 and K_2 are nontrivial knots, then $K_1 \# K_2$ is not $SU(2)$ -averse.*

Proof. We know that $K_1 \# K_2$ is a satellite $P(K_2)$, where $P(U) = K_1$. Indeed, we can remove a small open neighborhood of a meridian $m \subset \partial N(K_1)$ from S^3 to get a solid torus $S^1 \times D^2$ with embedded knot $P = K_1$, and this is the desired pattern. Similarly, there is a pattern P' such that $P'(K_1) = K_1 \# K_2$ and $P'(U) = K_2$. Proposition 10.2 applied to P and to P' now says that if $K_1 \# K_2$ is $SU(2)$ -averse, then each summand must be $SU(2)$ -averse as well and

$$r(K_1) = r(K_1 \# K_2) = r(K_2).$$

We let r denote the common limit slope, recalling that $r \neq 0$.

The knot group of $K_1 \# K_2$ is an amalgamated free product

$$\pi_1(S^3 \setminus K_1) *_Z \pi_1(S^3 \setminus K_2)$$

in which we identify the meridians $\mu_1 \sim \mu_2$, with peripheral data $\mu_{\#} = \mu_1 = \mu_2$ and $\lambda_{\#} = \lambda_1 \lambda_2$. Since both K_1 and K_2 are $SU(2)$ -averse, they satisfy case (2) of Theorem 4.1, so the pillowcase images of $X^*(K_1)$ and $X^*(K_2)$ contain arcs of the form

$$\beta = -r\alpha + c_i, \quad \frac{\pi}{2} \leq \alpha < \frac{\pi}{2} + \varepsilon$$

for some constants c_1 and c_2 respectively and some $\varepsilon > 0$. For each α in this interval, we let $\rho_{\alpha}^j : \pi_1(S^3 \setminus K_j) \rightarrow SU(2)$, $j = 1, 2$, be corresponding representations such that

$$\rho_{\alpha}^j(\mu_j) = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{-i\alpha} \end{pmatrix}, \quad \rho_{\alpha}^j(\lambda_j) = \begin{pmatrix} e^{i(-r\alpha+c_j)} & 0 \\ 0 & e^{-i(-r\alpha+c_j)} \end{pmatrix}.$$

Combining these produces a representation $\rho_\alpha^\# = \rho_\alpha^1 * \rho_\alpha^2$ of $\pi_1(S^3 \setminus (K_1 \# K_2))$, with

$$\rho_\alpha^\#(\mu_\#) = \begin{pmatrix} e^{i\alpha} & 0 \\ 0 & e^{-i\alpha} \end{pmatrix}, \quad \rho_\alpha^\#(\lambda_\#) = \begin{pmatrix} e^{i(-2r\alpha+c_1+c_2)} & 0 \\ 0 & e^{-i(-2r\alpha+c_1+c_2)} \end{pmatrix}.$$

But the $\rho_\alpha^\#$ produce a line segment of slope $-2r$ in the pillowcase image of $X^*(K_1 \# K_2)$, so by Theorem 3.5 we must actually have $r(K_1 \# K_2) = 2r$, and since $r \neq 2r$ this is impossible. \square

10.2. Cables. If p and q are relatively prime integers with $q \geq 2$, and K is an arbitrary knot, we define the (p, q) -cable $C_{p,q}(K)$ to be a peripheral curve in $\partial N(K)$ in the class $\mu^p \lambda^q$. This is clearly a satellite with winding number q , and if $|p| \geq 2$ then $C_{p,q}(U)$ is the (p, q) -torus knot.

Proposition 10.5. *If K is a nontrivial, $SU(2)$ -averse knot with limit slope $r(K) = \frac{p}{q}$, where p and q are relatively prime and $q \geq 2$, then the (p, q) -cable $C = C_{p,q}(K)$ is $SU(2)$ -averse with limit slope $r(C) = pq$.*

Proof. According to Theorem 8.2, there is an integer $N \geq 1$ such that $(\frac{p}{q} + \frac{1}{kN})$ -surgery on K is $SU(2)$ -averse for all but finitely many k . Gordon [Gor83, Corollary 7.3] showed that

$$S_{pq+\frac{1}{n}}^3(C_{p,q}(K)) \cong S_{\frac{p}{q}+\frac{1}{q^2n}}^3(K)$$

for all $n \geq 1$, and the latter is $SU(2)$ -cyclic for all but finitely many multiples n of N , so $C_{p,q}(K)$ has an infinite sequence of $SU(2)$ -cyclic slopes which converges to pq . \square

Theorem 10.6. *Let K be a nontrivial knot, and let p and q be relatively prime with $q \geq 2$. The cable $C_{p,q}(K)$ is $SU(2)$ -averse if and only if K is also $SU(2)$ -averse with limit slope $r(K) = \frac{p}{q}$, and in this case $r(C_{p,q}(K)) = pq$.*

Proof. One direction was already established in Proposition 10.5, so we will suppose that $C_{p,q}(K)$ is $SU(2)$ -averse. By Proposition 10.3, K is $SU(2)$ -averse with $r(C_{p,q}(K)) = q^2 r(K)$. If $p \neq \pm 1$ then $C_{p,q}(U) = T_{p,q}$ is not the unknot, so $r(C_{p,q}(K)) = r(T_{p,q}) = pq$ by Proposition 10.2 and we conclude that $r(K) = \frac{p}{q}$.

The only remaining cases are $p = \pm 1$, but we begin with some generalities about arbitrary cables. Many of the details below are adapted from the proof of [NZ17, Theorem 2.8]. Letting $C = C_{p,q}$ denote the corresponding pattern in $S^1 \times D^2$, the complement $(S^1 \times D^2) \setminus C$ is Seifert fibered over the annulus with one singular fiber of order q , where the generic fibers on $S^1 \times \partial D^2$ have slope

$$p[\mu_K] + q[\lambda_K] = p[\{\text{pt}\} \times \partial D^2] + q[S^1 \times \{\text{pt}\}],$$

and those on $\partial N(C)$ have slope $pq[\mu_C] + [\lambda_C]$; see [Gor83, Lemma 7.2]. These are central in $\pi_1((S^1 \times D^2) \setminus C)$, so any representation

$$\rho : \pi_1(S^3 \setminus C_{p,q}(K)) \rightarrow SU(2)$$

whose restriction to $\pi_1((S^1 \times D^2) \setminus C)$ is irreducible must send these to the center of $SU(2)$, meaning that $\rho(\mu_C^{pq} \lambda_C)$ is either I or $-I$.

We now assume that $p = \pm 1$ and let $\rho_K : \pi_1(S^3 \setminus K) \rightarrow SU(2)$ be an irreducible representation satisfying $\rho_K(\mu_K^{\pm 1} \lambda_K^q) = I$; this exists since $\pm \frac{1}{q}$ -surgery on K is not $SU(2)$ -cyclic, by [KM04a]. Then $\rho_K(\mu_K) \notin \{I, -I\}$, or else ρ_K would be reducible, and since

$\rho_K(\lambda_K^q) \notin \{I, -I\}$ as a result, the same is true of $\rho_K(\lambda_K)$. We will use ρ_K to construct a family of representations

$$\rho_s : \pi_1(S^3 \setminus C_{\pm 1, q}(K)) \rightarrow SU(2)$$

which are irreducible on $(S^1 \times D^2) \setminus C$, hence satisfy $\rho_s(\mu_C^{\pm q} \lambda_C) \in \{I, -I\}$, but for which the trace of $\rho_s(\mu_C)$ takes values everywhere in an open interval. This produces a line segment of slope $\mp q$ in the pillowcase image of $X^*(C_{\pm 1, q}(K))$, which implies that $C_{\pm 1, q}(K)$ has limit slope $\pm q$. But then we have $r(K) = \frac{1}{q^2} r(C_{\pm 1, q}(K)) = \pm \frac{1}{q}$ as above, and this contradicts $|r(K)| > 2$, so the proof will be complete.

Taking $C = C_{\pm 1, q} \subset S^1 \times D^2$, we have a presentation

$$\pi_1((S^1 \times D^2) \setminus C) = \langle h, \lambda_K \mid h^q \lambda_K = \lambda_K h^q \rangle$$

with $\mu_C = h\lambda_K$, as deduced in [NZ17]; here h is freely homotopic to the singular fiber of $(S^1 \times D^2) \setminus C$. (Note that we are using K to refer to the companion knot and C to refer to the cabling pattern, in contrast with [NZ17].) Thus we can extend ρ_K to a representation of $\pi_1(S^3 \setminus C_{\pm 1, q}(K))$ by finding an element of $SU(2)$ whose q th power commutes with $\rho_K(\lambda_K)$.

Viewing $SU(2)$ as the unit quaternions for convenience, we can write $\rho_K(\lambda_K) = e^{i\beta}$ up to conjugacy for some β , noting that $\sin(\beta) \neq 0$ since $e^{i\beta} \notin \{1, -1\}$. We define a family of purely imaginary unit quaternions by

$$v_s = \cos(s)i + \sin(s)j, \quad 0 \leq s \leq \pi.$$

Then for all s in the same range we extend ρ_K to $\rho_s : \pi_1(S^3 \setminus C_{\pm 1, q}(K)) \rightarrow SU(2)$ with pillowcase coordinates (α_s, β_s) by setting

$$\rho_s(h) = \cos\left(\frac{\pi}{q}\right) + \sin\left(\frac{\pi}{q}\right)v_s.$$

It is clear that $\rho_s(h^q) = -1$, which commutes with $\rho_s(\lambda_K) = e^{i\beta}$, and so these representations are well-defined. For $0 < s < \pi$ they restrict to irreducible representations on $\pi_1((S^1 \times D^2) \setminus C)$, since $\rho_s(h)$ does not commute with $e^{i\beta}$. From $\rho_s(\mu_K) = \rho_s(h)\rho_s(\lambda_K)$ we also compute that

$$\cos(\alpha_s) = \operatorname{Re}(\rho_s(\mu_K)) = \cos\left(\frac{\pi}{q}\right)\cos(\beta) - \cos(s)\sin\left(\frac{\pi}{q}\right)\sin(\beta).$$

Since $\sin(\frac{\pi}{q})\sin(\beta) \neq 0$, the value of $\cos(\alpha_s)$ and hence of α_s covers an open interval of values for $0 < s < \pi$, as desired. \square

Theorem 10.6 says that if any cable at all of K is $SU(2)$ -averse then exactly one cable $C_{p, q}(K)$ is, with limit slope $r(C_{p, q}(K)) = pq$; and in this case, not only is K also $SU(2)$ -averse but the limit slope $r(K) = \frac{p}{q}$ cannot be an integer. Since $SU(2)$ -averse small knots (which include torus knots) and cables must have integral limit slopes, we conclude the following.

Corollary 10.7. *Cables of small knots and of nontrivial cables are never $SU(2)$ -averse.*

We have the following special case, since algebraic knots are iterated cables of the unknot.

Corollary 10.8. *An algebraic knot is $SU(2)$ -averse if and only if it is a torus knot.*

10.3. Whitehead doubles. Using what we know so far about satellites, we cannot say much about whether the Whitehead double $Wh(K)$ of a nontrivial knot is $SU(2)$ -averse: neither Proposition 10.2 nor Proposition 10.3 can be used here, since $Wh(U)$ is the unknot and the corresponding pattern knot has winding number zero. On the other hand, we can completely answer Conjecture 1.5 for Whitehead doubles by other means.

Proposition 10.9. *If K is a nontrivial knot, then its Whitehead double $Wh(K)$ is not $SU(2)$ -averse.*

Proof. Since $Wh(K)$ has Seifert genus 1, its smooth slice genus is at most 1. If it is smoothly slice, then Theorem 9.3 says that it is not $SU(2)$ -averse. Otherwise its slice genus is exactly 1, and then since its Alexander polynomial is 1 we know by Proposition 9.4 that it is not $SU(2)$ -averse. \square

11. MONTESINOS KNOTS AND KNOTS WITH SMALL CROSSING NUMBER

In this section we provide evidence for Conjecture 1.5, that the only $SU(2)$ -averse knots are the torus knots, by proving it for several classes of examples. Our main tools will be:

- Theorem 7.3 for small knots, and in particular the inequality $\det(K) \leq |r(K)| - 1$;
- restrictions on the roots of the Alexander polynomial from Proposition 6.5; and
- Theorem 9.3 and Proposition 9.4, which assert that an $SU(2)$ -averse knot cannot be smoothly slice, and that if it has slice genus 1 then its Seifert genus is also 1.

We have obtained data about knots with low crossing number from the Knot Atlas [BNM⁺] and KnotInfo [CL], and we rely on SnapPy [CDGW] for a computation. In general, we only discuss each knot with the chirality specified in these tables, but if K is not $SU(2)$ -averse then neither is its mirror \overline{K} .

First, we will use Theorem 7.3 together with bounds on the boundary slopes of two-bridge knots, and of Montesinos knots with three rational tangles, in terms of the crossing number $c(K)$ to verify Conjecture 1.5 for these knots.

Theorem 11.1. *Let K be an $SU(2)$ -averse knot which is an alternating Montesinos knot with at most three rational tangles. Then K is a $(2, 2n + 1)$ -torus knot.*

Proof. By hypothesis, the set of finite boundary slopes of K has diameter equal to $2c(K)$; this is due to Mattman, Maybrun, and Robinson [MMR08] in the two-bridge case (i.e., at most two rational tangles), and to Ichihara and Mizushima [IM08] for alternating Montesinos knots in general. Both 0 and the limit slope $r(K)$ are boundary slopes, so we must have $|r(K)| \leq 2c(K)$. Two-bridge knots are small [HT85], as are Montesinos knots with at most three rational tangles [Oer84], so Theorem 7.3 implies that

$$\det(K) \leq |r(K)| - 1 \leq 2c(K) - 1. \quad (11.1)$$

Theorem 10.4 says that K is prime, so by Corollary 11.3 (which is stated and proved below), we see that K is either a $(2, 2n + 1)$ torus knot, a twist knot, 6_2 , or 7_3 . But we have already seen in Example 7.2 that the trefoil is the only nontrivial, $SU(2)$ -averse twist knot, so this leaves only 6_2 and 7_3 . These have Alexander polynomials

$$\begin{aligned} \Delta_{6_2}(t) &= -t^2 + 3t - 3 + 3t^{-1} - t^{-2} \\ \Delta_{7_3}(t) &= 2t^2 - 3t + 3 - 3t^{-1} + 2t^{-2}, \end{aligned}$$

both of which are irreducible and not cyclotomic; since $\Delta_{6_2}(-1) = -11$ and $\Delta_{7_3}(i) = -1$, Proposition 6.5 says that neither knot is $SU(2)$ -averse. \square

Corollary 11.3, which was used in the proof of Theorem 11.1, asserts that the only prime alternating knots for which $\det(K) \leq 2c(K)$ are torus knots, twist knots, 6_2 , and 7_3 . It is an easy consequence of the following proposition, which in turn is a slight refinement of a theorem of Crowell [Cro59, Theorem 6.5].

Proposition 11.2. *If K is a prime alternating knot, then $\det(K) \geq 3c(K) - 8$ unless K is a $(2, 2k + 1)$ torus knot or a twist knot.*

Proof. Crowell [Cro59, Theorem 6.5] proved that any prime alternating knot K satisfies $\det(K) \geq 2c(K) - 3$ unless K is of “elementary torus type”; in the latter case, its Alexander polynomial has the form

$$\Delta_K(t) \doteq 1 - t + t^2 - \dots + (-t)^n$$

[Cro59, (6.6)] for some n , which guarantees by [OS05, Proposition 4.1] that it is a $(2, n + 1)$ torus knot. We assume from now on that K is not a torus knot.

Let B be the black graph associated to a checkerboard coloring of a diagram of K with $c(K)$ crossings. We let b and w denote the number of vertices and faces of B , or equivalently the number of faces and vertices of the white graph. Then $b, w \geq 3$ by [Cro59, (6.3)], and since B is planar with $c(K)$ edges, we have $b + w = c(K) + 2$.

The proof of [Cro59, Theorem 6.5] combines the relation

$$(b - 1)(w - 1) = 2(c(K) - 2) + (b - 3)(w - 3),$$

which is stated as the inequality [Cro59, Theorem 6.4] since $(b - 3)(w - 3) \geq 0$, with the inequality

$$\det(K) - 1 \geq (b - 1)(w - 1) \tag{11.2}$$

of [Cro59, Theorem 5.10]. We note that $(b - 3)(w - 3)$ is equal to

$$(b - 3)(c(K) - b - 1) = \left(\frac{c(K)}{2} - 2\right)^2 - \left(b - 1 - \frac{c(K)}{2}\right)^2,$$

so if neither $b - 3$ nor $w - 3$ is zero, then it is minimized when $b - 3 = 1$ or $w - 3 = 1$. In particular, if $(b - 3)(w - 3)$ is nonzero then it is at least $c(K) - 5$, and hence either

$$\det(K) \geq 2c(K) - 3 + (b - 3)(w - 3) \geq 3c(K) - 8$$

or one of $b - 3$ and $w - 3$ is zero.

In the remaining cases, we take $b = 3$ without loss of generality; in this case B has three vertices, $c(K)$ edges, and $w = c(K) - 1$ faces. Crowell deduces the inequality (11.2) from the fact that for alternating knots, $\det(K)$ is the number of spanning trees of B , so we must determine the cases in which (given $b = 3$ and $w = c(K) - 1$) the graph B has at most $3c(K) - 9$ spanning trees.

Label the vertices of B as v_1, v_2, v_3 , and let e_i denote the number of edges between v_j and v_k whenever $\{i, j, k\} = \{1, 2, 3\}$; note that $e_i \geq 1$ for all i , since there are no vertices whose removal from B disconnects it (see [Cro59, (5.3) and (5.4)]). There are $e_1e_2 + e_2e_3 + e_3e_1$ spanning trees and $c(K) = e_1 + e_2 + e_3$, so we wish to solve the inequality

$$e_1e_2 + e_2e_3 + e_3e_1 \leq 3(e_1 + e_2 + e_3) - 9.$$

Reordering the v_i if needed so that $e_1 \leq e_2 \leq e_3$, we rearrange the inequality to get

$$(e_1 + e_2 - 3)e_3 + (e_1 - 3)(e_2 - 3) \leq 0.$$

If $e_1 + e_2 \geq 3$, then it follows that $e_2 \geq 2$ (hence $2e_2 - 3 > 0$) and that

$$(e_1 + e_2 - 3)e_2 + (e_1 - 3)(e_2 - 3) \leq 0,$$

or equivalently $e_2^2 \leq (3 - e_1)(2e_2 - 3)$. But this implies that $e_2^2 \leq 2(2e_2 - 3)$, which has no solutions, so we must therefore have $e_1 + e_2 < 3$, which means that $e_1 = e_2 = 1$.

We conclude that up to permuting the coordinates, the possible values of (e_1, e_2, e_3) are $(1, 1, a)$ for some $a \geq 4$. In each of these cases, the corresponding black graph can only come from a twist knot, completing the proof. \square

Corollary 11.3. *If K is a prime alternating knot, then $\det(K) > 2c(K)$ unless K is a $(2, 2k + 1)$ torus knot, a twist knot, 6_2 , or 7_3 .*

Proof. We inspect the knot types with up to 7 crossings by hand to see that the knots with $\det(K) \leq 2c(K)$ are either torus knots (3_1 , 5_1 , and 7_1), twist knots (4_1 , 5_2 , 6_1 , and 7_2), or 6_2 or 7_3 . Assuming that $c(K) > 7$ and K is not a torus knot or a twist knot, we have $\det(K) \geq 3c(K) - 8 > 2c(K) - 1$ by Proposition 11.2, and since $\det(K)$ is always odd we in fact have $\det(K) > 2c(K)$, so there are no further examples. \square

We now begin to verify Conjecture 1.5 for knots with low crossing number.

Theorem 11.4. *If a knot with crossing number at most 9 is $SU(2)$ -averse, then it is a torus knot.*

Proof. Most knots with at most 8 crossings are alternating Montesinos knots with at most three tangles, and thus handled by Theorem 11.1; the exceptions are 8_n for $16 \leq n \leq 21$, of which 8_{19} is the $(3, 4)$ torus knot. The remaining five knots all have slice genus at most 1 and Seifert genus at least 2, so Theorem 9.3 and Proposition 9.4 say that they are not $SU(2)$ -averse.

Similarly, there are 49 knot types with crossing number 9, and the knot 9_n is alternating and Montesinos with at most three rational tangles for $1 \leq n \leq 28$ and $n = 30, 31, 35, 36, 37$, so Theorem 11.1 applies for these values of n . The remaining knots are 9_n for n in

$$29, 32, 33, 34, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49.$$

Among these knots, all but 9_{38} , 9_{43} , and 9_{49} are either smoothly slice, or have smooth slice genus 1 but Seifert genus greater than 1, so they cannot be $SU(2)$ -averse by Theorem 9.3 and Proposition 9.4.

We now use Proposition 6.5 to rule out 9_{38} , since

$$\Delta_{9_{38}}(t) = (t - 1 + t^{-1})(5t - 9 + 5t^{-1})$$

and the roots of $5t - 9 + 5t^{-1}$ lie on the unit circle but are not roots of unity; and 9_{49} , since

$$\Delta_{9_{49}}(t) = 3t^2 - 6t + 7 - 6t^{-1} + 3t^{-2}$$

is irreducible with $\Delta_{9_{49}}(e^{i\pi/3}) = -2$. This leaves only the small Montesinos knot 9_{43} : if it were $SU(2)$ -averse then $r(9_{43})$ would be one of its boundary slopes, which are

$$-4, 0, 6, 8, 32/3$$

by [HO89, Dun01], but since $\det(9_{43}) = 13$, it cannot satisfy $\det(9_{43}) < |r(9_{43})|$. \square

Theorem 11.5. *The only 10-crossing knot which is $SU(2)$ -averse is $10_{124} = T(3, 5)$, with the possible exception of 10_{98} .*

Proof. The 10-crossing knots are labeled 10_n for $n \leq 165$, and 10_n is alternating and Montesinos with at most three rational tangles for all $n \leq 78$, so Theorem 11.1 applies to them. For $79 \leq n \leq 165$, the knot 10_n is either smoothly slice, or has slice genus 1 and Seifert genus at least 2, for all n except 124 (i.e., the $(3, 5)$ torus knot) and

$$80, 85, 92, 98, 100, 101, 111, 120, 127, 128, 134, \\ 139, 142, 145, 149, 150, 152, 154, 157, 160, 161.$$

Thus Theorem 9.3 and Proposition 9.4 rule out all of the remaining 10-crossing knots except for these.

Among the remaining 21 knot types 10_n , most of these are not $SU(2)$ -averse by Proposition 6.5. The Alexander polynomials of the following knots have no cyclotomic factors, and are negative at the indicated points on the unit circle:

$$\Delta_{10_n}(e^{i\pi/4}) < 0: \quad n = 80, 92, 101, 111, 127, 128, 134, 145, 149, 150, 154, 157, 160, 161 \\ \Delta_{10_n}(e^{i\pi/6}) < 0: \quad n = 120, 152.$$

Proposition 6.5 can also be applied to the Alexander polynomials

$$\Delta_{10_{85}}(t) = (t - 1 + t^{-1})(t^3 - 3t^2 + 4t - 3 + 4t^{-1} - 3t^{-2} + t^{-3}), \\ \Delta_{10_{139}}(t) = (t - 1 + t^{-1})(t^3 - t + 1 - t^{-1} + t^{-3}), \\ \Delta_{10_{142}}(t) = (t - 1 + t^{-1})(2t^2 - t - 1 - t^{-1} + 2t^{-2}).$$

Indeed, if we call the second factors $f_{85}(t)$, $f_{139}(t)$, and $f_{142}(t)$ respectively, then these are real-valued on the unit circle and equal to 1 at $t = 1$, with $f_{85}(-1) = -19$, $f_{139}(e^{i\pi/3}) = -2$, and $f_{142}(i) = -5$, so that each $f_n(t)$ and hence $\Delta_{10_n}(t)$ has a simple root on the unit circle which is not a root of unity.

This leaves only the knots 10_{98} and 10_{100} . Since 10_{100} is small, we use SnapPy [CDGW] to compute the list of boundary slopes of spun normal surfaces in its complement:

$$\text{Manifold('10_100').normal_boundary_slopes()}$$

The list includes all nonzero boundary slopes since 10_{100} is a hyperbolic knot in S^3 , and thus it gives us an upper bound of 20 on the absolute value of a boundary slope for 10_{100} . This upper bound is smaller than $\det(10_{100}) = 65$, so Theorem 7.3 says that it cannot be $SU(2)$ -averse. \square

For 10_{98} , Kitano and Suzuki [KS08] showed that its knot group admits a surjection onto the knot group of 3_1 (or possibly its mirror) which sends $\mu_{10_{98}}$ to μ_{3_1} and $\lambda_{10_{98}}$ to $\lambda_{3_1}^2$. Arguing as in the proof of Proposition 10.2, we see that every $SU(2)$ -cyclic slope for 10_{98} is twice an $SU(2)$ -cyclic slope for 3_1 , and since 3_1 has limit slope ± 6 we must have $r(10_{98}) = \pm 12$ if 10_{98} is indeed $SU(2)$ -cyclic.

We include the sign ambiguity in the above discussion because we have not checked the chirality of the groups given in [KS08]. One can distinguish between 10_{98} and its mirror $\overline{10_{98}}$ as follows: since these are alternating knots, any Legendrian representative of either satisfies the inequality

$$tb(K) + |r(K)| \leq -\sigma(K) - 1$$

[Pla04, Corollary 1], where tb and r are the Thurston-Bennequin and rotation numbers of K , and where the right-handed trefoil has signature $\sigma(3_1) = -2$ by convention. One of these knot types has a Legendrian representative with $tb(K) = 1$ (though existing knot tables do not seem to agree on whether this knot is 10_{98} or $\overline{10_{98}}$), and so it must be the

one with signature -4 , which is 10_{98} . The proof of [BS16, Theorem 1.10] now shows that no negative surgery on this knot can ever be an instanton L-space, so if it is $SU(2)$ -averse then its limit slope would have to be positive by Theorem 9.1, and hence $r(10_{98}) = 12$.

APPENDIX A. C^r -APPROXIMATION THROUGH SHEARING MAPS

In this appendix, we prove Theorem 4.5, following the strategy of the proof of [Zen16, Theorem 3.3].

We start by fixing some notational and linguistic conventions which are self-suggesting by the fact that the tangent bundle of the torus is parallelizable. If X is a vector field on T^2 we regard it as a map $T^2 \rightarrow \mathbb{R}^2$. We consider its derivative as a map $DX : T^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$, and if the point p on the torus is fixed we denote by DX_p the corresponding linear map. The expression $\|DX\|_\infty$ denotes the supremum of the norms $\|DX_p\|$ as p varies over T^2 .

Likewise we regard the flow ϕ_X^t as a map $T^2 \rightarrow \mathbb{R}^2$, and we consider the derivative as a map $D\phi_X^t : T^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

By a slight abuse of notation, we shall denote by

$$\|\zeta - \xi\|_{C^r} := \sup_{(x,y) \in T^2} d(\zeta(x,y), \xi(x,y)) + \sum_{l=1}^r \left\| D^{(l)}\zeta - D^{(l)}\xi \right\|_\infty$$

the C^r -distance of the maps $\zeta, \xi : T^2 \rightarrow T^2$.

The Fourier decomposition. The first point is that Lemma 3.4 in [Zen16] generalizes to the C^l norm for any $l \geq 1$ if the vector field X one starts with is differentiable infinitely often. More precisely, for any smooth vector field X , and for any $\varepsilon > 0$, there is a sum of Fourier vector fields $Z = W_1 + \dots + W_m$ such that we have $\|X - Z\|_{C^l} < \varepsilon$. A Fourier vector field W_i is of the form

$$W_i(x, y) = \mathbf{a}_i \sin(\mathbf{k}_i \cdot (x, y)) + \mathbf{b}_i \cos(\mathbf{k}_i \cdot (x, y)),$$

where $\mathbf{a}_i, \mathbf{b}_i \in \mathbb{R}^2$, and $\mathbf{k}_i \in \mathbb{Z}^2$. It is shown in [Zen16, Lemma 3.4] that if the vector field Z is divergence-free, then the flow of the vector fields W_i are isotopies through shearing maps.

A few lemmata. In the sequel we denote by \mathbb{R}_+ the non-negative real integers. The following lemma is a generalization of [Zen16, Lemma 3.6].

Lemma A.1. *Let $r \geq 0$. For any $0 \leq l \leq r$, there are functions*

$$a_l : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$$

which are continuous and monotonically increasing in both variables, such that the following holds. Let $X : T^2 \rightarrow \mathbb{R}^2$ be a vector field on T^2 with flow $\phi_X^t : T^2 \rightarrow T^2$, and suppose that for any $l \leq r + 1$ its derivatives satisfy

$$\left\| D^{(i)}X \right\|_\infty \leq K_i \quad \text{for all } i \leq l$$

for some positive constants K_i . Then:

(1) *For any $p, q \in T^2$, we have*

$$\left\| \phi_X^t(p) - \phi_X^t(q) \right\| \leq a_0(t, K_1) \cdot \|p - q\|.$$

(2) *For any $p, q \in T^2$ and any $l \geq 1$, we have*

$$\left\| (D^{(l)}\phi_X^t)(p) - (D^{(l)}\phi_X^t)(q) \right\| \leq t \cdot a_l(t, K_{l+1}) \cdot \|p - q\|.$$

Proof. The case $r = 0$ is Lemma 3.6 in [Zen16], with $a_0(t, K_1) = e^{K_1 t}$. For $r \geq 1$, the proof will be by induction on r .

By definition the flow ϕ_X^t satisfies the differential equation

$$\frac{d\phi_X^t}{dt}(p) = X(\phi_X^t(p)),$$

for any $p \in T^2$, which integrated gives

$$\phi_X^t(p) = p + \int_0^t X(\phi_X^s(p)) ds. \quad (\text{A.1})$$

Differentiating this equation with respect to p yields

$$D\phi_X^t(p) = \text{id} + \int_0^t DX_{\phi_X^s(p)} \circ D\phi_X^s(p) ds. \quad (\text{A.2})$$

Therefore we get, using the triangle inequality and the mean value theorem,

$$\begin{aligned} \|D\phi_X^t(p) - D\phi_X^t(q)\| &\leq \int_0^t \left\| DX_{\phi_X^s(p)} \circ D\phi_X^s(p) - DX_{\phi_X^s(q)} \circ D\phi_X^s(q) \right\| ds \\ &\leq \int_0^t \left\| DX_{\phi_X^s(p)} \circ D\phi_X^s(p) - DX_{\phi_X^s(q)} \circ D\phi_X^s(p) \right\| ds \\ &\quad + \int_0^t \left\| DX_{\phi_X^s(q)} \circ D\phi_X^s(p) - DX_{\phi_X^s(q)} \circ D\phi_X^s(q) \right\| ds \\ &\leq \|D\phi_X^s\|_{L^\infty(T^2 \times [0, t])} \cdot K_2 \cdot \int_0^t \|\phi_X^s(p) - \phi_X^s(q)\| ds \\ &\quad + \|DX\|_\infty \int_0^t \|D\phi_X^s(p) - D\phi_X^s(q)\| ds. \end{aligned}$$

But [Zen16, Lemma 3.6] states that $\|\phi_X^s(p) - \phi_X^s(q)\| \leq e^{K_1 s} \|p - q\|$. Therefore, we get the inequality

$$\begin{aligned} \|D\phi_X^t(p) - D\phi_X^t(q)\| &\leq \|D\phi_X^s\|_{L^\infty(T^2 \times [0, t])} \cdot K_2 \cdot t \cdot e^{K_1 t} \|p - q\| \\ &\quad + \|DX\|_\infty \int_0^t \|D\phi_X^s(p) - D\phi_X^s(q)\| ds. \end{aligned}$$

Gronwall's inequality now yields

$$\|D\phi_X^t(p) - D\phi_X^t(q)\| \leq t \cdot \|p - q\| \cdot \|D\phi_X^s\|_{L^\infty(T^2 \times [0, t])} \cdot K_2 \cdot \exp(t(\|DX\|_\infty + K_1)) \quad (\text{A.3})$$

for all $p, q \in T^2$, and for all $t \geq 0$.

In a similar but easier application of Gronwall's inequality, Equation (A.2) yields

$$\|D\phi_X^s\|_{L^\infty(T^2 \times [0, t])} \leq \exp(K_1 t). \quad (\text{A.4})$$

Hence the bound in Equation (A.3) implies the $l = 1$ case of the lemma once we set $a_1(t, K_2) := K_2 \cdot \exp(3K_2 t)$.

We suppose now that the result holds for $r - 1$ for some $r \geq 2$. Differentiating Equation (A.2), we get the formula

$$\begin{aligned} (D^{(r)}\phi_X^t)(p) &= \int_0^t L^{(r-1)}(s) ds \\ &+ \int_0^t (DX)_{\phi_X^s(p)} \circ (D^{(r)}\phi_X^s)(p) ds. \end{aligned} \tag{A.5}$$

Here, the term $L^{(r-1)}(s)$ is a polynomial expression of derivatives of X and of ϕ_X^s of order up to $r - 1$. A precise expression could be given, but we don't need this here. The formula for higher derivatives of a composition of functions of one variable is known as Faà di Bruno's formula.

The claim now follows easily from this, using the same sort of estimates as above, the induction step, and then Gronwall's inequality to obtain an inequality for $\|D^{(r)}\phi_X^s\|_\infty$, and for the difference $\|(D^{(r)}\phi_X^s)(p) - (D^{(r)}\phi_X^s)(q)\|$. This will use the fact that sums and products of non-negative monotonely increasing functions are again monotonely increasing. We leave the details as an exercise for the interested reader. \square

One can prove the following generalization of [Zen16, Lemma 3.7] along the same lines.

Lemma A.2. *Let $r \geq 0$. Then for any $l \leq r$ there are functions*

$$b_l: \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$$

which are continuous and monotonically increasing in all three variables, such that the following holds:

- (1) *For all $t \geq 0$ and $K \geq 0$ we have $b_l(t, 0, K) = 0$.*
- (2) *Let $X, Y: T^2 \rightarrow \mathbb{R}^2$ be vector fields on T^2 , and suppose that for any $l \leq r + 1$ the derivatives of X satisfy*

$$\|D^{(i)}X\|_\infty \leq K_l \quad \text{for all } i \leq l$$

for some positive constants K_l . Then the flows $\phi_X^t, \phi_Y^t: T^2 \rightarrow T^2$ satisfy

$$\|(D^{(l)}\phi_X^t)(p) - (D^{(l)}\phi_Y^t)(p)\| \leq t \cdot b_l(t, \|X - Y\|_{C^l}, K_{l+1}).$$

for any $p \in T^2$.

Proof. The case $r = 0$ is Lemma 3.7 of [Zen16], with $b_0(t, d, K_1) = d \cdot e^{K_1 t}$. The proof for $r \geq 1$ again goes by induction. To start the induction, let $r = 1$. Applying the triangle inequality as above, one gets

$$\begin{aligned} \|D\phi_X^t(p) - D\phi_Y^t(p)\| &\leq \int_0^t \|DX_{\phi_X^s(p)} \circ D\phi_X^s(p) - DY_{\phi_Y^s(p)} \circ D\phi_Y^s(p)\| ds \\ &\leq \int_0^t \|DX_{\phi_X^s(p)} \circ D\phi_X^s(p) - DX_{\phi_Y^s(p)} \circ D\phi_X^s(p)\| ds \\ &\quad + \int_0^t \|DX_{\phi_Y^s(p)} \circ D\phi_X^s(p) - DX_{\phi_Y^s(p)} \circ D\phi_Y^s(p)\| ds \\ &\quad + \int_0^t \|DX_{\phi_Y^s(p)} \circ D\phi_Y^s(p) - DY_{\phi_Y^s(p)} \circ D\phi_Y^s(p)\| ds \end{aligned}$$

$$\begin{aligned}
 &\leq \|D\phi_X^s\|_{L^\infty(T^2 \times [0,t])} \cdot K_2 \cdot \int_0^t \|\phi_X^s(p) - \phi_Y^s(p)\| \, ds \\
 &\quad + \|DX\|_\infty \int_0^t \|D\phi_X^s(p) - D\phi_Y^s(p)\| \, ds \\
 &\quad + t \|DX - DY\|_\infty \cdot \|D\phi_Y^s\|_{L^\infty(T^2 \times [0,t])}.
 \end{aligned}$$

Now [Zen16, Lemma 3.7] yields the estimate $\|\phi_X^s(p) - \phi_Y^s(p)\| \leq s \|X - Y\|_\infty e^{K_1 s}$, where K_1 can be taken as a Lipschitz constant for X . Gronwall's inequality now yields

$$\begin{aligned}
 \|D\phi_X^t(p) - D\phi_Y^t(p)\| &\leq t (\|DX - DY\|_\infty \cdot \|D\phi_Y^s\|_{L^\infty(T^2 \times [0,t])} \\
 &\quad + K_2 \cdot \|X - Y\|_\infty \cdot e^{K_1 t} \cdot \|D\phi_X^s\|_{L^\infty(T^2 \times [0,t])}) \cdot \exp(t \|DX\|_\infty).
 \end{aligned}$$

This has the desired form once we notice that by (A.4) and the triangle inequality,

$$\begin{aligned}
 \|D\phi_Y^s\|_\infty &\leq \exp(\|DY\|_\infty s) \leq \exp(\|DX - DY\|_\infty s + \|DX\|_\infty s) \\
 &\leq \exp(\|DX - DY\|_\infty s + K_1 s).
 \end{aligned}$$

Assuming the claim holds up to $r - 1$, the induction step now follows again from Equation (A.5) similarly as in the proof of the previous lemma. Again, we leave the details to the reader. \square

We will also need a generalization of [Zen16, Lemma 3.11].

Lemma A.3. *Let $r \geq 0$. Then for any $l \leq r$ and any $m \geq 2$ there are functions*

$$c_l^{(m)} : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$$

which are continuous and monotonically increasing in all three variables, such that the following holds:

Let Z be a vector field given as a sum of vector fields $Z = W_0 + \dots + W_{m-1}$. Suppose that there are positive constants K_0, \dots, K_{r+1} and M_0, \dots, M_r such that for any $l \leq r + 1$ one has

$$\left\| D^{(i)}(W_{j_1} + \dots + W_{j_n}) \right\|_\infty \leq K_l$$

for all $i \leq l$ and all nonempty subsets $\{j_1, \dots, j_n\}$ of $\{0, \dots, m-1\}$, and that for any $l \leq r$ and $1 \leq n \leq m-1$ we have

$$\left\| D^{(i)}[W_n, W_0 + \dots + W_{n-1}] \right\|_\infty \leq M_l$$

for all $i \leq l$. Then for any $p \in T^2$ we have

$$\left\| (D^{(l)}\phi_Z^t)(p) - (D^{(l)}(\phi_{W_m}^t \circ \dots \circ \phi_{W_1}^t))(p) \right\| \leq t^2 \cdot c_l^{(m)}(t, M_l, K_{l+1})$$

for all $t \geq 0$.

Proof. We recall that for two vector fields X, Y the Lie bracket is defined to be

$$[X, Y](p) = \lim_{h \rightarrow 0} \frac{D\phi_Y^h(X(\phi_Y^{-h}(p)) - X(p))}{h} = \frac{d}{dt} [(\phi_Y^t)_*(X)](p)|_{t=0}.$$

Hence we have for any point p the equality

$$D\phi_Y^t(X(p)) = X(\phi_Y^t(p)) + t[X, Y](\phi_Y^t(p)) + R_{X,Y}(\phi_Y^t(p), t), \quad (\text{A.6})$$

where the term $R_{X,Y}$ satisfies

$$\lim_{t \rightarrow 0} \frac{R_{X,Y}(p, t)}{t} = 0$$

and this convergence is uniform in p . Moreover, any derivatives of R with respect to p have the same limit as t goes to 0, since we can interchange the order of the derivatives.

The proof of the lemma now goes by induction on l . The case $l = 0$ is the statement of [Zen16, Lemma 3.11]. We will prove the lemma for $l = 1$, leaving the induction to higher l to the interested reader. This case will itself proceed by induction on m , the number of summands in the expression $Z = W_0 + \cdots + W_{m-1}$, beginning with $m = 2$.

Suppose we have $Z = W_0 + W_1$. Differentiating the composition $\phi_{W_1}^t \circ \phi_{W_0}^t$ and using the above remark, we obtain

$$\begin{aligned} \frac{d}{dt} \phi_{W_1}^t(\phi_{W_0}^t(p)) &= \frac{d\phi_{W_1}^t}{dt}(\phi_{W_0}^t(p)) + (\phi_{W_1}^t)_* \left(\frac{d\phi_{W_0}^t}{dt}(p) \right) \\ &= W_1(\phi_{W_1}^t(\phi_{W_0}^t(p))) + (\phi_{W_1}^t)_* W_0(\phi_{W_0}^t(p)) \\ &= (W_1 + W_0)(\phi_{W_1}^t(\phi_{W_0}^t(p))) \\ &\quad + t[W_0, W_1](\phi_{W_1}^t(\phi_{W_0}^t(p))) + R_{W_0, W_1}(\phi_{W_1}^t(\phi_{W_0}^t(p)), t). \end{aligned} \tag{A.7}$$

Integrating this equation, and differentiating with respect to p , we obtain the integral equality

$$\begin{aligned} D(\phi_{W_1}^t \circ \phi_{W_0}^t)(p) &= \text{id} + \int_0^t D(W_1 + W_0) \circ D(\phi_{W_1}^s \circ \phi_{W_0}^s)(p) ds \\ &\quad + \int_0^t s D([W_1, W_0]) \circ D(\phi_{W_1}^s \circ \phi_{W_0}^s)(p) ds \\ &\quad + \int_0^t s D\left(\frac{R}{s}\right)(\phi_{W_1}^s(\phi_{W_0}^s(p))) ds. \end{aligned} \tag{A.8}$$

We subtract this from the corresponding integral equality which is satisfied by the derivative of the flow $\phi_{W_1+W_0}^t$, as in (A.2), and this yields the estimate

$$\begin{aligned} &\| (D\phi_{W_1+W_0}^t)(p) - D(\phi_{W_1}^t \circ \phi_{W_0}^t)(p) \| \\ &\leq \int_0^t \left\| D(W_1 + W_0)_{\phi_{W_1+W_0}^s(p)} \circ D\phi_{W_1+W_0}^s(p) \right. \\ &\quad \left. - D(W_1 + W_0)_{(\phi_{W_1}^s \circ \phi_{W_0}^s)(p)} \circ D(\phi_{W_1}^s \circ \phi_{W_0}^s)(p) \right\| ds \\ &\quad + \int_0^t s \| [W_1, W_0] \|_{C^1} \| D(\phi_{W_1}^s \circ \phi_{W_0}^s) \| ds \\ &\quad + \int_0^t s \left\| D\frac{R}{s} \right\| ds \end{aligned}$$

$$\begin{aligned}
&\leq \int_0^t \left\| D(W_1 + W_0)_{\phi_{W_1+W_0}^s(p)} \right\| \left\| D\phi_{W_1+W_0}^s(p) - D(\phi_{W_1}^s \circ \phi_{W_0}^s)(p) \right\| ds \\
&\quad + \int_0^t \left\| D(W_1 + W_0)_{\phi_{W_1+W_0}^s(p)} - D(W_1 + W_0)_{(\phi_{W_1}^s \circ \phi_{W_0}^s)(p)} \right\| \left\| D(\phi_{W_1}^s \circ \phi_{W_0}^s)(p) \right\| ds \\
&\quad + \frac{t^2}{2} \max_{s \in [0,t]} (\| [W_1, W_0] \|_{C^1} \cdot \| D(\phi_{W_1}^s \circ \phi_{W_0}^s) \|) \\
&\quad + \frac{t^2}{2} \max_{s \in [0,t]} \left\| \frac{R}{s} \right\|_{C^1} \\
&\leq K_1 \int_0^t \left\| D\phi_{W_1+W_0}^s(p) - D(\phi_{W_1}^s \circ \phi_{W_0}^s)(p) \right\| ds \\
&\quad + K_2 \int_0^t \left\| \phi_{W_1+W_0}^s(p) - (\phi_{W_1}^s \circ \phi_{W_0}^s)(p) \right\| \left\| D(\phi_{W_1}^s \circ \phi_{W_0}^s)(p) \right\| ds \\
&\quad + \frac{t^2}{2} \max_{s \in [0,t]} (\| [W_1, W_0] \|_{C^1} \cdot \| D(\phi_{W_1}^s \circ \phi_{W_0}^s) \|) \\
&\quad + \frac{t^2}{2} \max_{s \in [0,t]} \left\| \frac{R}{s} \right\|_{C^1}. \tag{A.9}
\end{aligned}$$

By the induction hypothesis we have

$$\left\| \phi_{W_1+W_0}^s - \phi_{W_1}^s \circ \phi_{W_0}^s \right\| \leq s^2 c_0^{(2)}(s, M_0, K_1)$$

for all s . Furthermore one easily shows that

$$\left\| D(\phi_{W_1}^s \circ \phi_{W_0}^s) \right\|_{\infty} \leq e^{K_1 s}$$

holds for all s . If we plug these two inequalities into the estimate (A.9), Gronwall's inequality yields the desired result for $l = 1$; here we define

$$c_1^{(2)}(t, M, K) := (K c_0^{(2)}(t, M, K) e^{Kt} + \frac{1}{2} M e^{Kt} + \frac{1}{2} r_1(t, M, K)) e^{Kt},$$

where $r_1(t, M, K)$ is a function that can be explicitly described in terms of R_{W_1, W_0} , which was defined in Equation (A.6) above. We leave the induction in the cases $m \geq 3$ as an exercise for the interested reader. \square

We are now ready to start with the proof of Theorem 4.5.

Strategy of the proof. We observe that an isotopy $(\psi_t)_{t \in [0,1]}$ as in the statement of Theorem 4.5 is the flow of a time-dependent, divergence-free vector field X_t satisfying

$$\frac{d\psi^t(p)}{dt} = X_t(\psi^t(p)) \tag{A.10}$$

for any $p \in T^2$ and any $t \in [0, 1]$. Integrating the vector field from time t_0 to time t sends a point $p \in T^2$ to $\psi^t((\psi^{t_0})^{-1}(p))$.

Let $\varepsilon > 0$ be given, and suppose that we want to approximate ψ_t to within a distance of at most ε in the C^r norm. The proof of Theorem 4.5 is an approximation in three steps, just as in the proof of [Zen16, Theorem 3.3].

- (1) We approximate the flow ψ_t of X_t by a composition of flows of time-independent vector fields $X_i := X_{i/n}$, $i = 0, \dots, n-1$. More precisely, we define the isotopies $(\Theta_{(X_j)}^t) : T^2 \rightarrow T^2$ on each interval $\frac{i}{n} \leq t \leq \frac{i+1}{n}$ by

$$\Theta_{(X_j)}^t := \phi_{X_i}^{t-i/n} \circ \phi_{X_{i-1}}^{1/n} \circ \dots \circ \phi_{X_0}^{1/n}. \quad (\text{A.11})$$

We will show that by taking n large enough, we get

$$\left\| \Theta_{(X_j)}^t - \psi^t \right\|_{C^r} < \frac{\varepsilon}{3}$$

for all $t \in [0, 1]$.

- (2) We approximate each of the X_i by a finite sum Z_i of Fourier vector fields, using the C^r -version of [Zen16, Lemma 3.4] mentioned above. We define $\Theta_{(Z_j)}^t$ to be the analogue of (A.11) above with Z_i in place of X_i , and by keeping track of the accumulated error, we find that

$$\left\| \Theta_{(X_j)}^t - \Theta_{(Z_j)}^t \right\|_{C^r} < \frac{\varepsilon}{3}$$

for all $t \in [0, 1]$, assuming each Z_j was chosen sufficiently C^r -close to X_j .

- (3) We write each Z_j as a finite Fourier series

$$Z_j = \sum_{r=0}^{m_j-1} W_r^{(j)},$$

where each $W_r^{(j)}$ is a shearing vector field in some direction in \mathbb{Z}^2 . We will approximate the flow of Z_j , which occurs over a time interval of length at most $\frac{1}{n}$ in $\Theta_{(Z_j)}^t$, by successive flows along each of the summands $W_r^{(j)}$.

For each Z_j , we fix some $k_j \in \mathbb{N}$ and define an isotopy $\Xi_{(W_r^{(j)})}^t$ as follows. For $0 \leq t \leq \frac{1}{k_j m_j n}$, we flow along $W_0^{(j)}$ with speed m_j ; then we flow along $W_1^{(j)}$ with speed m_j for $\frac{1}{k_j m_j n} \leq t \leq \frac{2}{k_j m_j n}$; and so on, until we flow along $W_{m_j-1}^{(j)}$ during the m_j th interval of length $\frac{1}{k_j m_j n}$ and a total time of $\frac{1}{k_j n}$ has elapsed. We then repeat this k_j times to get the desired $\Xi_{(W_r^{(j)})}^t$, defined for $0 \leq t \leq \frac{1}{n}$.

We now approximate $\Theta_{(Z_j)}^t$ by an isotopy $\Omega_{(Z_j)}^t$, defined in terms of $\Xi_{(W_r^{(j)})}^t$ by

$$\Omega_{(Z_j)}^t := \Xi_{(W_r^{(i)})}^{t-i/n} \circ \Xi_{(W_r^{(i-1)})}^{1/n} \circ \dots \circ \Xi_{(W_r^{(0)})}^{1/n} \quad (\text{A.12})$$

for $\frac{i}{n} \leq t \leq \frac{i+1}{n}$, where $i = 0, \dots, n-1$. We will show that if for each sum $Z_j = \sum_{r=0}^{m_j-1} W_r^{(j)}$ we choose the corresponding k_j large enough, then

$$\left\| \Theta_{(Z_j)}^t - \Omega_{(Z_j)}^t \right\|_{C^r} < \frac{\varepsilon}{3}$$

for all $t \in [0, 1]$.

First step. We start by approximating the isotopy ψ_t by $\Theta_{(X_j)}^t$, as defined in (A.11). Thus we need an estimate that bounds the distance from ψ^t to $\Theta_{(X_j)}^t$ in the C^r -topology.

Lemma A.4. *Let $r \geq 0$. Then for any $l \leq r$ and $n \in \mathbb{N}$, there are functions*

$$h_l^{(n)} : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$$

which are continuous and monotonically increasing in all three variables, such that the following holds:

(1) *The functions $h_l^{(n)}$ have the property that*

$$\lim_{x \rightarrow 0^+} h_l^{(n)}(t, x, K) = 0,$$

and the convergence is uniform in n , and also in t and K taken from compact subsets of \mathbb{R}_+ .

(2) *Suppose that for any $l \leq r + 1$ we have*

$$\left\| D^{(i)} X_t \right\|_\infty \leq K_l$$

for all $i \leq l$ and all $t \geq 0$. Then we have the estimate

$$\left\| D^{(l)} \psi^t - D^{(l)} \Theta_{(X_j)}^t \right\|_\infty \leq h_l^{(n)} \left(t, \max_{j=0, \dots, n-1} \max_{\frac{j}{n} \leq t \leq \frac{j+1}{n}} \|X_j - X_t\|_{C^1(T^2)}, K_{l+1} \right)$$

for all $t \in [0, 1]$.

Proof. We prove the lemma by induction on l . The case $l = 0$ is the conclusion of [Zen16, Lemma 3.9]. In fact, we can define the function $h_0^{(n)}$ by the formula

$$h_0^{(n)}(t, x, K) := \left(t - \frac{j}{n} \right) x e^{K(t - \frac{j}{n})} + \frac{x}{n} \sum_{i=0}^{j-1} e^{K(t - \frac{i}{n})}$$

where $\frac{j}{n} \leq t \leq \frac{j+1}{n}$. This function clearly is bounded uniformly by $f_0(t, x, K) := t x e^{Kt}$, and hence the claim follows.

For simplicity of notation and more clarity of the argument, we will treat the case $l = 1$ only, and we will leave the induction argument for $l \geq 2$ to the interested reader.

The proof of Lemma A.2 extends verbatim to the case of vector fields which depend on time. In our case, we will compare the flow of X_t to that of X_0 . The conclusion we get is the uniform bound

$$\left\| D\psi^t - D\phi_{X_0}^t \right\| \leq t \cdot b_1(t, \sup_{s \in [0, t]} \|X_0 - X_s\|_{C^1(T^2)}, K_2) \tag{A.13}$$

which holds for all $t \geq 0$. It is also straightforward to verify the bounds

$$\left\| D\psi^t \right\|_\infty \leq e^{K_1 t} \quad \text{and} \quad \left\| D\phi_{X_j}^t \right\|_\infty \leq e^{K_1 t} \tag{A.14}$$

for any $j = 0, \dots, n - 1$, cf. Equation (A.4).

We claim that for $\frac{j}{n} \leq t \leq \frac{j+1}{n}$ we have

$$\left\| (D\psi^t)(p) - (D\Theta_{(X_j)}^t)(p) \right\|$$

$$\begin{aligned}
&\leq \left(t - \frac{j}{n}\right) \left[b_1 \left(\frac{1}{n}, \sup_{s \in [\frac{j}{n}, \frac{j+1}{n}]} \|X_j - X_s\|_{C^1(T^2)}, K_2 \right) \right. \\
&\quad \left. + a_1 \left(\frac{1}{n}, K_2 \right) h_0^{(n)} \left(\frac{j}{n}, \sup_{s \in [\frac{j}{n}, \frac{j+1}{n}]} \|X_i - X_s\|_{C^1(T^2)}, K_1 \right) \right] e^{K_1 \frac{j}{n}} \\
&\quad + \sum_{i=0}^{j-1} \frac{1}{n} \left[b_1 \left(\frac{1}{n}, \sup_{s \in [\frac{i}{n}, \frac{i+1}{n}]} \|X_j - X_s\|_{C^1(T^2)}, K_2 \right) \right. \\
&\quad \left. + a_1 \left(\frac{1}{n}, K_2 \right) h_0^{(n)} \left(\frac{i}{n}, \max_{i=0, \dots, n-1} \sup_{s \in [\frac{i}{n}, \frac{i+1}{n}]} \|X_i - X_s\|_{C^1(T^2)}, K_1 \right) \right] e^{K_1 \frac{i}{n}}.
\end{aligned} \tag{A.15}$$

Here a_1 and b_1 are some functions satisfying the statement of Lemmas A.1 and A.2 above.

We prove this claim by induction on j . The case $j = 0$, and hence the claim for times $0 \leq t \leq \frac{1}{n}$ follows immediately from (A.13). (Notice that $h_0^{(n)}(0, x, K) = 0$.)

Suppose the claim holds for $t \leq \frac{j}{n}$. Using the triangle inequality, Lemmas A.1 and A.2, and Equation (A.14) above, we obtain for $\frac{j}{n} \leq t \leq \frac{j+1}{n}$ the bound

$$\begin{aligned}
\left\| (D\psi^t)(p) - (D\Theta_{(X_j)}^t)(p) \right\| &= \left\| D(\psi^t \circ (\psi^{\frac{j}{n}})^{-1})_{\psi^{\frac{j}{n}}(p)} \circ (D\psi^{\frac{j}{n}})(p) - (D\phi_{X_j}^{t-\frac{j}{n}})_{\Theta^{\frac{j}{n}}(p)} \circ D\Theta_{(X_j)}^{\frac{j}{n}}(p) \right\| \\
&\leq \left\| D(\psi^t \circ (\psi^{\frac{j}{n}})^{-1})_{\psi^{\frac{j}{n}}(p)} - (D\phi_{X_j}^{t-\frac{j}{n}})_{\psi^{\frac{j}{n}}(p)} \right\| \cdot \left\| (D\psi^{\frac{j}{n}})(p) \right\| \\
&\quad + \left\| (D\phi_{X_j}^{t-\frac{j}{n}})_{\psi^{\frac{j}{n}}(p)} - (D\phi_{X_j}^{t-\frac{j}{n}})_{\Theta^{\frac{j}{n}}(p)} \right\| \cdot \left\| (D\psi^{\frac{j}{n}})(p) \right\| \\
&\quad + \left\| (D\psi^{\frac{j}{n}})(p) - (D\Theta_{(X_j)}^{\frac{j}{n}})(p) \right\| \cdot \left\| (D\phi_{X_j}^{t-\frac{j}{n}})_{\Theta^{\frac{j}{n}}(p)} \right\| \\
&\leq \left(t - \frac{j}{n}\right) b_1 \left(t - \frac{j}{n}, \sup_{s \in [\frac{j}{n}, t]} \|X_j - X_s\|_{C^1(T^2)}, K_2 \right) e^{K_1 \frac{j}{n}} \\
&\quad + \left(t - \frac{j}{n}\right) a_1 \left(t - \frac{j}{n}, K_2 \right) \left\| \psi^{\frac{j}{n}}(p) - \Theta^{\frac{j}{n}}(p) \right\| e^{K_1 \frac{j}{n}} \\
&\quad + \left\| (D\psi^{\frac{j}{n}})(p) - (D\Theta_{(X_j)}^{\frac{j}{n}})(p) \right\| e^{K_1(t-\frac{j}{n})}.
\end{aligned}$$

Using the induction hypothesis to bound the last term, and the case $l = 0$ to bound the middle term, we clearly obtain the bound (A.15). Hence we define the function $h_1^{(n)}$ by the formula

$$\begin{aligned}
h_1^{(n)}(t, x, K) &:= \left(t - \frac{j}{n}\right) \left(b_1 \left(\frac{1}{n}, x, K \right) + a_1 \left(\frac{1}{n}, K \right) h_0^{(n)} \left(\frac{j}{n}, x, K \right) \right) e^{Kt} \\
&\quad + \sum_{i=0}^{j-1} \frac{1}{n} \left(b_1 \left(\frac{1}{n}, x, K \right) + a_1 \left(\frac{1}{n}, K \right) h_0^{(n)} \left(\frac{i}{n}, x, K \right) \right) e^{K \frac{i}{n}}
\end{aligned}$$

for $\frac{j}{n} \leq t \leq \frac{j+1}{n}$. The bound (A.15) implies statement (2) of the lemma for the case $l = 1$.

The functions $h_1^{(n)}$ satisfy the bound

$$h_1^{(n)}(t, x, K) \leq f_1(t, x, K) := (b_1(1, x, K) + a_1(1, K)f_0(t, x, K))e^{Kt}$$

for all $t \in [0, 1]$, where we recall that $f_0(t, x, K) = txe^{Kt}$. By the properties of the functions f_0 , a_1 and b_1 we clearly have

$$\lim_{x \rightarrow 0^+} f_1(t, x, K) = 0,$$

and this limit is uniform in t and K taken from compact subsets. This implies statement (1) of the lemma for the case $l = 1$. \square

The first statement of Lemma A.4 tells us that there are functions

$$f_l : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$$

such that

$$h_l^{(n)}(t, x, K_{r+1}) \leq f_l(x, K_{r+1})$$

for all $n \in \mathbb{N}$ and all $t \in [0, 1]$. Furthermore, these functions satisfy

$$\lim_{x \rightarrow 0^+} f_l(x, K) = 0$$

for all K , so we can choose $M > 0$ so small that $f_l(M, K_{r+1}) < \frac{\varepsilon}{3(r+1)}$ for all $l = 0, \dots, r$.

We now notice that the time dependent vector field $(X_t) : T^2 \times [0, 1] \rightarrow \mathbb{R}^2$ is equicontinuous, and all of its derivatives are also equicontinuous maps. In particular, given M as above, there is some $\delta > 0$ such that we have

$$\|X_s - X_t\|_{C^r(T^2)} < M \quad \text{whenever} \quad |s - t| < \delta.$$

Choosing n large enough so that $\frac{1}{n} < \delta$, the second conclusion of Lemma A.4 implies that

$$\left\| \Theta_{(X_j)}^t - \psi^t \right\|_{C^r(T^2)} < \frac{\varepsilon}{3}.$$

for all $t \in [0, 1]$, as desired.

Second step. We now approximate $\Theta_{(X_j)}^t$ by $\Theta_{(Z_j)}^t$, where each Z_j is a finite Fourier series which C^r -approximates X_j . The following is nearly a duplicate of Lemma A.4, with $\Theta_{(X_j)}^t$ and $\Theta_{(Z_j)}^t$ in place of ψ^t and $\Theta_{(X_j)}^t$.

Lemma A.5. *Let $r \geq 0$. Then for any $l \leq r$ and $n \in \mathbb{N}$, there are functions*

$$g_l^{(n)} : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$$

which are continuous and monotonically increasing in all three variables, such that:

(1) *The functions $g_l^{(n)}$ have the property that*

$$\lim_{x \rightarrow 0^+} g_l^{(n)}(t, x, K) = 0,$$

and the convergence is uniform in n , and also in t and K taken from compact subsets of \mathbb{R}_+ .

(2) Suppose that for any $l \leq r + 1$ we have

$$\left\| D^{(i)} X_j \right\|_{\infty} \leq K_l$$

for all $i \leq l$. Then we have the estimate

$$\left\| D^{(l)} \Theta_{(X_j)}^t - D^{(l)} \Theta_{(Z_j)}^t \right\|_{\infty} \leq g_l^{(n)}(t, \max_{j=0, \dots, n-1} \max_{\frac{j}{n} \leq t \leq \frac{j+1}{n}} \|X_j - Z_j\|_{C^l(T^2)}, K_{l+1})$$

for all $t \in [0, 1]$.

Proof. We repeat the proof of Lemma A.4 verbatim, using the same functions $h_l^{(n)}$ as before. The only details which need to be established first are bounds for each $l \leq r + 1$ of the form $\|D^{(i)} Z_j\|_{\infty} \leq K_l$, for all $i \leq l$ and all $j = 0, \dots, n - 1$. But clearly we have

$$\left\| D^{(i)} Z_j \right\|_{\infty} \leq \left\| D^{(i)} X_j \right\|_{\infty} + \left\| D^{(i)} (Z_j - X_j) \right\|_{\infty} \leq K_l + \|Z_j - X_j\|_{C^l},$$

and we can absorb the extra $\|Z_j - X_j\|_{C^l}$ in the definition of the functions $g_l^{(n)}$. \square

The first statement of the preceding lemma tells us that

$$\lim_{x \rightarrow 0^+} g_l^{(n)}(t, x, K_{r+1}) = 0$$

uniformly in $t \in [0, 1]$. (We don't need the statement about uniform convergence in n , as n was already fixed in the first approximation step.) We choose $M > 0$ small enough that $g_l^{(n)}(1, M, K_{r+1}) \leq \frac{\varepsilon}{3(r+1)}$ for all $l = 0, \dots, r$. Having fixed M , we choose the finite Fourier sums $Z_j = W_0^{(j)} + \dots + W_{m_j-1}^{(j)}$ so that we have

$$\|X_j - Z_j\|_{C^r} \leq M$$

for $j = 0, \dots, n - 1$. Now the second conclusion of Lemma A.5 implies that

$$\left\| \Theta_{(X_j)}^t - \Theta_{(Z_j)}^t \right\|_{C^r(T^2)} < \frac{\varepsilon}{3}$$

for all $t \in [0, 1]$.

Third step. We start with a lemma that describes the C^l -distance between the flow ϕ_Z^t of $Z = W_0 + \dots + W_{m-1}$ and the isotopy $\Xi_{(W_r)}^t$, defined as in step (3). To be explicit, suppose that we have fixed positive integers n and k , so that for any $t \in [0, \frac{1}{n}]$ we can find integers i, r such that

$$\frac{1}{n} \left(\frac{i}{k} + \frac{r}{km} \right) \leq t \leq \frac{1}{n} \left(\frac{i}{k} + \frac{r+1}{km} \right)$$

with $0 \leq i \leq k - 1$ and $0 \leq r \leq m - 1$. Then we define $\Xi_{(W_r)}^t$ by the formula

$$\Xi_{(W_r)}^t := \phi_{W_r}^{m(t - \frac{im+r}{kmn})} \circ \phi_{W_{r-1}}^{\frac{1}{kn}} \circ \dots \circ \phi_{W_0}^{\frac{1}{kn}} \circ \left(\phi_{W_{m-1}}^{\frac{1}{kn}} \circ \dots \circ \phi_{W_0}^{\frac{1}{kn}} \right)^i.$$

Lemma A.6. *Let $r \geq 0$. Then for any $l \leq r$ and $k \in \mathbb{N}$, there are functions*

$$d_l^{(k)} : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$$

which are continuous and monotonically increasing in all three variables, such that:

(1) The functions $d_l^{(k)}$ have the property that

$$\lim_{k \rightarrow \infty} d_l^{(k)}(t, M, K) = 0$$

for any M, K , and t .

(2) Let Z be a vector field given as a sum of vector fields $Z = W_0 + \cdots + W_{m-1}$. Suppose that there are positive constants K_0, \dots, K_{r+1} and M_0, \dots, M_r such that for any $l \leq r+1$ we have

$$\left\| D^{(i)}(W_{j_1} + \cdots + W_{j_n}) \right\|_{\infty} \leq K_l$$

for all $i \leq l$ and all finite subsets $\{j_1, \dots, j_n\}$ of $\{0, \dots, m-1\}$, and such that

$$\left\| D^{(i)}[W_n, W_0 + \cdots + W_{n-1}] \right\|_{\infty} \leq M_l$$

for all $i \leq l \leq r$ and all $n \geq 1$. Then we have the estimate

$$\left\| D^{(l)}\phi_Z^t - D^{(l)}\Xi_{(W_r)}^t \right\|_{\infty} \leq d_l^{(k)}(t, M_l, K_{l+1})$$

for all $t \geq 0$. (We recall here that $\Xi_{(W_r)}^t$ depends on k .)

Proof. The case $l = 0$ is the content of [Zen16, Lemma 3.12]. Again, we will show the case $l = 1$ and leave the induction for higher l to the interested reader; we start by noting that

$$\left\| D\Xi_{(W_r)}^t \right\|_{\infty} \leq e^{K_1 t} \tag{A.16}$$

for all $t \geq 0$.

We first claim that at times $t = \frac{j}{nk}$, where j is an integer, we have

$$\begin{aligned} \left\| D\Xi_{(W_r)}^{\frac{j}{nk}} - D\phi_Z^{\frac{j}{nk}} \right\|_{\infty} &\leq \sum_{i=0}^{j-1} \frac{1}{nk} \left[\frac{1}{nk} c_1 \left(\frac{1}{nk}, M_1, K_2 \right) \right. \\ &\quad \left. + a_1 \left(\frac{1}{nk}, K_2 \right) d_0^{(k)} \left(\frac{j}{nk}, M_1, K_2 \right) \right] e^{K_1 \frac{j}{nk}}. \end{aligned} \tag{A.17}$$

Here c_1 denotes a function $c_1^{(m)} : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ as provided by Lemma A.3 above; we omit the superscript (m) for convenience, as well as the (W_r) subscript on Ξ in the sequel.

The proof of (A.17) is by induction, analogous to the proof of the bound (A.15) in Lemma A.5; in this situation, we make use of Lemma A.3 above. By using the triangle inequality, followed by Lemma A.1 and (A.4), we obtain the estimate

$$\begin{aligned} \left\| (D\Xi^t)(p) - (D\phi_Z^t)(p) \right\| &= \left\| D(\Xi^{t-\frac{j}{nk}})_{\Xi_{\frac{j}{nk}}(p)} \circ (D\Xi_{\frac{j}{nk}})(p) - (D\phi_Z^{t-\frac{j}{nk}})_{\phi_Z^{\frac{j}{nk}}(p)} \circ D\phi_Z^{\frac{j}{nk}}(p) \right\| \\ &\leq \left\| D(\Xi^{t-\frac{j}{nk}})_{\Xi_{\frac{j}{nk}}(p)} - (D\phi_Z^{t-\frac{j}{nk}})_{\Xi_{\frac{j}{nk}}(p)} \right\| \cdot \left\| (D\Xi_{\frac{j}{nk}})(p) \right\| \\ &\quad + \left\| (D\phi_Z^{t-\frac{j}{nk}})_{\Xi_{\frac{j}{nk}}(p)} - (D\phi_Z^{t-\frac{j}{nk}})_{\phi_Z^{\frac{j}{nk}}(p)} \right\| \cdot \left\| (D\Xi_{\frac{j}{nk}})(p) \right\| \\ &\quad + \left\| (D\Xi_{\frac{j}{nk}})(p) - (D\phi_Z^{\frac{j}{nk}})(p) \right\| \cdot \left\| (D\phi_Z^{t-\frac{j}{nk}})_{\Theta_{\frac{j}{nk}}(p)} \right\| \end{aligned}$$

$$\begin{aligned}
&\leq \left\| D(\Xi^{t-\frac{j}{nk}})_{\Xi^{\frac{j}{nk}}(p)} - (D\phi_Z^{t-\frac{j}{nk}})_{\Xi^{\frac{j}{nk}}(p)} \right\| e^{K_1 \frac{j}{nk}} \\
&\quad + \left(t - \frac{j}{nk} \right) a_1 \left(t - \frac{j}{nk}, K_2 \right) \left\| \Xi^{\frac{j}{nk}}(p) - \phi_Z^{\frac{j}{nk}}(p) \right\| e^{K_1 \frac{j}{nk}} \\
&\quad + \left\| (D\Xi^{\frac{j}{nk}})(p) - (D\phi_Z^{\frac{j}{nk}})(p) \right\| e^{K_1(t-\frac{j}{nk})}. \tag{A.18}
\end{aligned}$$

Supposing we are trying to prove (A.17) for $t = \frac{j+1}{nk}$, the last two terms of (A.18) can be bounded using the case $l = 0$ of the lemma and the case $t = \frac{j}{nk}$ of (A.17) respectively. At time $t = \frac{j+1}{nk}$ the map $\Xi^{t-\frac{j}{nk}}$ is equal to $\Xi^{\frac{1}{nk}} = \phi_{W_{m-1}}^{\frac{1}{nk}} \circ \dots \circ \phi_{W_0}^{\frac{1}{nk}}$, and hence Lemma A.3 gives us the bound

$$\begin{aligned}
\left\| D(\Xi^{t-\frac{j}{nk}})_{\Xi^{\frac{j}{nk}}(p)} - (D\phi_Z^{t-\frac{j}{nk}})_{\Xi^{\frac{j}{nk}}(p)} \right\| &= \left\| D(\Xi^{\frac{1}{nk}})_{\Xi^{\frac{j}{nk}}(p)} - (D\phi_Z^{\frac{1}{nk}})_{\Xi^{\frac{j}{nk}}(p)} \right\| \\
&\leq \left(\frac{1}{nk} \right)^2 c_1 \left(\frac{1}{nk}, M_1, K_2 \right).
\end{aligned}$$

This bound together with (A.18) establishes the bound (A.17) by induction.

Now the intermediate times $\frac{j}{kn} \leq t \leq \frac{j+1}{kn}$ need a separate treatment, because for these times $\Xi^{t-\frac{j}{nk}}$ is not of the form where Lemma A.3 applies. In fact, for these times we can only establish a coarser estimate:

$$\begin{aligned}
\left\| D\Xi^{t-\frac{j}{nk}} - D\phi_Z^{t-\frac{j}{nk}} \right\|_{\infty} &\leq \left\| D\Xi^{t-\frac{j}{nk}} - \text{id} \right\|_{\infty} + \left\| D\phi_Z^{t-\frac{j}{nk}} - \text{id} \right\|_{\infty} \\
&\leq \left(t - \frac{j}{nk} \right) K_1 e^{K_1(t-\frac{j}{nk})} + \left(t - \frac{j}{nk} \right) K_1 e^{K_1(t-\frac{j}{nk})} \\
&= 2 \left(t - \frac{j}{nk} \right) K_1 e^{K_1(t-\frac{j}{nk})}. \tag{A.19}
\end{aligned}$$

The estimate $\left\| D\phi_Z^t - \text{id} \right\|_{\infty} \leq t K_1 e^{K_1 t}$ is an immediate application of Gronwall's inequality to the derivative of the flow equation of Z , i.e. equation (A.2), and we leave the estimate

$$\left\| D\Xi_Z^{t-\frac{j}{nk}} - \text{id} \right\|_{\infty} \leq \left(t - \frac{j}{nk} \right) K_1 e^{K_1(t-\frac{j}{nk})}$$

for $\frac{j}{kn} \leq t \leq \frac{j+1}{kn}$ as an exercise to the interested reader.

In total, we have now proved that for any $\frac{j}{nk} \leq t \leq \frac{j+1}{nk}$ we have

$$\begin{aligned}
\left\| D\Xi_{(W_r)}^t - D\phi_Z^t \right\|_{\infty} &\leq \sum_{i=0}^j \frac{1}{nk} \left[\max \left\{ \frac{1}{nk} c_1 \left(\frac{1}{nk}, M_1, K_2 \right), \frac{2}{nk} K_1 e^{\frac{K_1}{nk}} \right\} \right. \\
&\quad \left. + a_1 \left(\frac{1}{nk}, K_2 \right) d_0^{(k)} \left(\frac{j}{nk}, M_1, K_2 \right) \right] e^{K_1 \frac{i}{nk}}. \tag{A.20}
\end{aligned}$$

Now notice that for $\frac{j}{nk} \leq t < \frac{j+1}{nk}$ the sum on the right hand side has $j+1 = \lfloor nkt \rfloor + 1$ summands. Thus we clearly obtain the estimate

$$\begin{aligned} \left\| D\Xi_{(W_r)}^t - D\phi_Z^t \right\|_\infty &\leq 2 \left(t + \frac{1}{k} \right) \left(\max \left\{ \frac{1}{k} c_1(1, M_1, K_2), \frac{2}{k} K_1 e^{\frac{K_1}{k}} \right\} \right. \\ &\quad \left. + a_1 \left(\frac{1}{k}, K_2 \right) d_0^{(k)}(1, M_1, K_2) \right) e^{K_1 t}. \end{aligned} \quad (\text{A.21})$$

(The functions a_1 , c_1 , and $d_0^{(k)}$ are increasing in t , so we can thus obtain expressions independent of n .) We define $d_1^{(k)}(t, M_1, K_2)$ to be the right hand side of (A.21), and then

$$\lim_{k \rightarrow \infty} d_1^{(k)}(t, M, K) = 0$$

follows immediately from the fact that $\lim_{k \rightarrow \infty} d_0^{(k)}(t, M, K) = 0$. \square

We are now ready to compare $\Theta_{(Z_j)}^t$ to the map $\Omega_{(Z_j)}^t$ defined in equation (A.12) above. Recall that the definition of $\Omega_{(Z_j)}^t$ depends on the summands $W_0^{(j)}, \dots, W_{m_j-1}^{(j)}$ in $Z_j = W_0^{(j)} + \dots + W_{m_j-1}^{(j)}$ and on the integers k_j which appear in the definitions of the maps $\Xi_{(W_r^{(j)})}^t$, for $j = 0, 1, \dots, n-1$. Here n has been fixed in the first approximation step, and the vector fields Z_0, \dots, Z_{n-1} have been fixed in the second approximation step.

Lemma A.7. *Let $r \geq 0$. Then for any $l \leq r$ there are functions*

$$f_l^{(k_0, \dots, k_{n-1})} : \mathbb{R}_+ \times \mathbb{R}_+^n \times \mathbb{R}_+^n \rightarrow \mathbb{R},$$

which are continuous and monotonically increasing in all variables, such that:

(1) For any tuples $\mathbf{M} = (M_0, \dots, M_{n-1})$ and $\mathbf{K} = (K_0, \dots, K_{n-1})$ we have

$$\lim_{k_{\min} \rightarrow \infty} f_l^{(k_0, \dots, k_{n-1})}(t, \mathbf{M}, \mathbf{K}) = 0,$$

where k_{\min} is defined as the minimum of the numbers k_0, \dots, k_{n-1} .

(2) Let Z_j be a vector field of the form $W_0^{(j)} + \dots + W_{m_j-1}^{(j)}$ for $0 \leq j \leq n-1$. Suppose that for each j there are positive constants $K_0^{(j)}, \dots, K_{r+1}^{(j)}$ and $M_0^{(j)}, \dots, M_r^{(j)}$ such that for any $l \leq r+1$ one has

$$\left\| D^{(i)}(W_{i_1}^{(j)} + \dots + W_{i_q}^{(j)}) \right\|_\infty \leq K_l^{(j)}$$

for all $i \leq l$ and all subsets $\{i_1, \dots, i_q\}$ of $\{0, \dots, m_j-1\}$, and

$$\left\| D^{(i)}[W_q^{(j)}, W_0^{(j)} + \dots + W_{q-1}^{(j)}] \right\|_\infty \leq M_l^{(j)}$$

for all $i \leq l \leq r$ and $1 \leq q \leq m_j-1$. Then we have the estimate

$$\left\| D^{(l)}\Theta_{(Z_j)}^t - D^{(l)}\Omega_{(W_r^{(j)})}^t \right\|_\infty \leq f_l^{(k_0, \dots, k_{n-1})}(t, (M_l^{(0)}, \dots, M_l^{(n-1)}), (K_{l+1}^{(0)}, \dots, K_{l+1}^{(n-1)}))$$

for all $t \geq 0$.

Proof. The theme of the proof is by now familiar. The case $l = 0$ is the statement of [Zen16, Lemma 3.13]. For the case $l = 1$, and for times $\frac{j}{n} \leq t \leq \frac{j+1}{n}$, one starts with the estimate

$$\begin{aligned} \left\| D\Omega_{(Z_j)}^t - D\Theta_{(Z_j)}^t \right\| &= \left\| D\Xi_{(W_r^{(j)})}^{t-\frac{j}{n}} \circ D\Omega_{(Z_j)}^{\frac{j}{n}} - D\phi_{Z_j}^{t-\frac{j}{n}} \circ D\Theta_{(Z_j)}^{\frac{j}{n}} \right\| \\ &\leq \left\| D\Omega_{(Z_j)}^{t-\frac{j}{n}} - D\phi_{Z_j}^{t-\frac{j}{n}} \right\| \left\| D\Omega_{(Z_j)}^{\frac{j}{n}} \right\| + \left\| D\phi_{Z_j}^{t-\frac{j}{n}} \right\| \left\| D\Omega_{(Z_j)}^{\frac{j}{n}} - D\Theta_{(Z_j)}^{\frac{j}{n}} \right\|. \end{aligned} \tag{A.22}$$

Now it is simple to verify that $\left\| D\Omega_{(Z_j)}^{\frac{j}{n}} \right\| \leq e^{\frac{1}{n}(K_1^{(0)} + \dots + K_1^{(j-1)})}$, and together with Lemma A.6 this gives inductively for $\frac{j}{n} \leq t \leq \frac{j+1}{n}$ the estimate

$$\begin{aligned} \left\| D\Omega_{(Z_j)}^t - D\Theta_{(Z_j)}^t \right\| &\leq \sum_{i=0}^{j-1} d_1^{(k_i)} \left(\frac{1}{n}, M_1^{(j)}, K_2^{(j)} \right) e^{\frac{1}{n}(K_1^{(0)} + \dots + K_1^{(i-1)})} \\ &\quad + d_1^{(k_j)} \left(t - \frac{j}{n}, M_1^{(j)}, K_2^{(j)} \right) e^{\frac{1}{n}(K_1^{(0)} + \dots + K_1^{(j-1)})}, \end{aligned}$$

where the $d_1^{(k)}$ are functions satisfying the conclusion of Lemma A.6. The right hand side defines the function $f_1^{(k_0, \dots, k_{n-1})}$, and the fact that

$$\lim_{k_{\min} \rightarrow \infty} f_l^{(k_0, \dots, k_{n-1})}(t, \mathbf{M}, \mathbf{K}) = 0$$

follows from the first conclusion of Lemma A.6.

We leave the induction steps for $l \geq 2$ as an exercise for the interested reader. \square

Once the vector fields Z_j and their decomposition as sums $Z_j = W_0^{(j)} + \dots + W_{m_j-1}^{(j)}$ are fixed, there are constants $K_l^{(j)}$ and $M_l^{(j)}$ satisfying the hypotheses of Lemma A.7. Lemma A.7 then allows us to choose the multi-index (k_0, \dots, k_{n-1}) large enough so that we have

$$f_l^{(k_0, \dots, k_{n-1})}(t, (M_l^{(0)}, \dots, M_l^{(n-1)}), (K_{l+1}^{(0)}, \dots, K_{l+1}^{(n-1)})) < \frac{\varepsilon}{3(r+1)}$$

for all $l = 0, \dots, r$, and it follows that

$$\left\| \Theta_{(Z_j)}^t - \Omega_{(Z_j)}^t \right\|_{C^r} < \frac{\varepsilon}{3}$$

for all $t \in [0, 1]$. This concludes the proof of Theorem 4.5.

REFERENCES

- [BCR98] J. Bochnak, M. Coste, and M.-F. Roy. *Real algebraic geometry*, volume 36 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1998. Translated from the 1987 French original, Revised by the authors.
- [BD95] P. J. Braam and S. K. Donaldson. Floer's work on instanton homology, knots and surgery. In *The Floer memorial volume*, volume 133 of *Progr. Math.*, pages 195–256. Birkhäuser, Basel, 1995.
- [BNM⁺] D. Bar-Natan, S. Morrison, et al. The Knot Atlas. <http://katlas.org>. August 23, 2017.
- [BS16] J. A. Baldwin and S. Sivek. Stein fillings and $SU(2)$ representations. arXiv:1611.05629, 2016.
- [Bur90] G. Burde. $SU(2)$ -representation spaces for two-bridge knot groups. *Math. Ann.*, 288(1):103–119, 1990.
- [BZ98] S. Boyer and X. Zhang. On Culler-Shalen seminorms and Dehn filling. *Ann. of Math. (2)*, 148(3):737–801, 1998.

- [CCG⁺94] D. Cooper, M. Culler, H. Gillet, D. D. Long, and P. B. Shalen. Plane curves associated to character varieties of 3-manifolds. *Invent. Math.*, 118(1):47–84, 1994.
- [CDGW] M. Culler, N. M. Dunfield, M. Goerner, and J. R. Weeks. SnapPy, a computer program for studying the geometry and topology of 3-manifolds. Available at <http://snappy.computop.org> (04/04/2016).
- [CGLS87] M. Culler, C. M. Gordon, J. Luecke, and P. B. Shalen. Dehn surgery on knots. *Ann. of Math. (2)*, 125(2):237–300, 1987.
- [CL] J. C. Cha and C. Livingston. KnotInfo: Table of Knot Invariants. <http://www.indiana.edu/~knotinfo>. August 23, 2017.
- [CL96] D. Cooper and D. D. Long. Remarks on the A -polynomial of a knot. *J. Knot Theory Ramifications*, 5(5):609–628, 1996.
- [Cro59] R. H. Crowell. Nonalternating links. *Illinois J. Math.*, 3:101–120, 1959.
- [CS83] M. Culler and P. B. Shalen. Varieties of group representations and splittings of 3-manifolds. *Ann. of Math. (2)*, 117(1):109–146, 1983.
- [CS84] M. Culler and P. B. Shalen. Bounded, separating, incompressible surfaces in knot manifolds. *Invent. Math.*, 75(3):537–545, 1984.
- [CS99] O. Collin and B. Steer. Instanton Floer homology for knots via 3-orbifolds. *J. Differential Geom.*, 51(1):149–202, 1999.
- [DG04] N. M. Dunfield and S. Garoufalidis. Non-triviality of the A -polynomial for knots in S^3 . *Algebr. Geom. Topol.*, 4:1145–1153 (electronic), 2004.
- [Don02] S. K. Donaldson. *Floer homology groups in Yang-Mills theory*, volume 147 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2002. With the assistance of M. Furuta and D. Kotschick.
- [Dun01] N. M. Dunfield. A table of boundary slopes of Montesinos knots. *Topology*, 40(2):309–315, 2001.
- [Flo88] A. Floer. An instanton-invariant for 3-manifolds. *Comm. Math. Phys.*, 118(2):215–240, 1988.
- [Flo90] A. Floer. Instanton homology, surgery, and knots. In *Geometry of low-dimensional manifolds, 1 (Durham, 1989)*, volume 150 of *London Math. Soc. Lecture Note Ser.*, pages 97–114. Cambridge Univ. Press, Cambridge, 1990.
- [FM66] R. H. Fox and J. W. Milnor. Singularities of 2-spheres in 4-space and cobordism of knots. *Osaka J. Math.*, 3:257–267, 1966.
- [Gol84] W. M. Goldman. The symplectic nature of fundamental groups of surfaces. *Adv. in Math.*, 54(2):200–225, 1984.
- [Gor83] C. M. Gordon. Dehn surgery and satellite knots. *Trans. Amer. Math. Soc.*, 275(2):687–708, 1983.
- [Hat82] A. E. Hatcher. On the boundary curves of incompressible surfaces. *Pacific J. Math.*, 99(2):373–377, 1982.
- [Hed10] M. Hedden. Notions of positivity and the Ozsváth-Szabó concordance invariant. *J. Knot Theory Ramifications*, 19(5):617–629, 2010.
- [Her94] C. M. Herald. Legendrian cobordism and Chern-Simons theory on 3-manifolds with boundary. *Comm. Anal. Geom.*, 2(3):337–413, 1994.
- [Her97] C. M. Herald. Flat connections, the Alexander invariant, and Casson’s invariant. *Comm. Anal. Geom.*, 5(1):93–120, 1997.
- [HK98] M. Heusener and J. Kroll. Deforming abelian SU(2)-representations of knot groups. *Comment. Math. Helv.*, 73(3):480–498, 1998.
- [HO89] A. Hatcher and U. Oertel. Boundary slopes for Montesinos knots. *Topology*, 28(4):453–480, 1989.
- [HT85] A. Hatcher and W. Thurston. Incompressible surfaces in 2-bridge knot complements. *Invent. Math.*, 79(2):225–246, 1985.
- [IM08] K. Ichihara and S. Mizushima. Crossing number and diameter of boundary slope set of Montesinos knot. *Comm. Anal. Geom.*, 16(3):565–589, 2008.
- [Kla91] E. P. Klassen. Representations of knot groups in SU(2). *Trans. Amer. Math. Soc.*, 326(2):795–828, 1991.
- [KM95] P. B. Kronheimer and T. S. Mrowka. Gauge theory for embedded surfaces. II. *Topology*, 34(1):37–97, 1995.
- [KM04a] P. B. Kronheimer and T. S. Mrowka. Dehn surgery, the fundamental group and SU(2). *Math. Res. Lett.*, 11(5-6):741–754, 2004.

- [KM04b] P. B. Kronheimer and T. S. Mrowka. Witten’s conjecture and property P. *Geom. Topol.*, 8:295–310 (electronic), 2004.
- [KM10a] P. Kronheimer and T. Mrowka. Instanton Floer homology and the Alexander polynomial. *Algebr. Geom. Topol.*, 10(3):1715–1738, 2010.
- [KM10b] P. Kronheimer and T. Mrowka. Knots, sutures, and excision. *J. Differential Geom.*, 84(2):301–364, 2010.
- [KM11] P. B. Kronheimer and T. S. Mrowka. Khovanov homology is an unknot-detector. *Publ. Math. Inst. Hautes Études Sci.*, (113):97–208, 2011.
- [KS08] T. Kitano and M. Suzuki. A partial order in the knot table. II. *Acta Math. Sin. (Engl. Ser.)*, 24(11):1801–1816, 2008.
- [Lan66] S. Lang. *Introduction to transcendental numbers*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1966.
- [Lim10] Y. Lim. Instanton homology and the Alexander polynomial. *Proc. Amer. Math. Soc.*, 138(10):3759–3768, 2010.
- [Lin16] J. Lin. $SU(2)$ -cyclic surgeries on knots. *Int. Math. Res. Not. IMRN*, (19):6018–6033, 2016.
- [MMR94] J. W. Morgan, T. Mrowka, and D. Ruberman. *The L^2 -moduli space and a vanishing theorem for Donaldson polynomial invariants*. Monographs in Geometry and Topology, II. International Press, Cambridge, MA, 1994.
- [MMR08] T. W. Mattman, G. Maybrun, and K. Robinson. 2-bridge knot boundary slopes: diameter and genus. *Osaka J. Math.*, 45(2):471–489, 2008.
- [Mos71] L. Moser. Elementary surgery along a torus knot. *Pacific J. Math.*, 38:737–745, 1971.
- [Ni07] Y. Ni. Knot Floer homology detects fibred knots. *Invent. Math.*, 170(3):577–608, 2007.
- [NZ17] Y. Ni and X. Zhang. Detection of knots and a cabling formula for A -polynomials. *Algebr. Geom. Topol.*, 17(1):65–109, 2017.
- [Oer84] U. Oertel. Closed incompressible surfaces in complements of star links. *Pacific J. Math.*, 111(1):209–230, 1984.
- [OS05] P. Ozsváth and Z. Szabó. On knot Floer homology and lens space surgeries. *Topology*, 44(6):1281–1300, 2005.
- [Pla04] O. Plamenevskaya. Bounds for the Thurston-Bennequin number from Floer homology. *Algebr. Geom. Topol.*, 4:399–406, 2004.
- [Sca15] C. W. Scaduto. Instantons and odd Khovanov homology. *J. Topol.*, 8(3):744–810, 2015.
- [Smy15] C. Smyth. Seventy years of Salem numbers. *Bull. Lond. Math. Soc.*, 47(3):379–395, 2015.
- [SW06] D. S. Silver and W. Whitten. Knot group epimorphisms. *J. Knot Theory Ramifications*, 15(2):153–166, 2006.
- [Tau90] C. H. Taubes. Casson’s invariant and gauge theory. *J. Differential Geom.*, 31(2):547–599, 1990.
- [Zen16] R. Zentner. Integer homology 3-spheres admit irreducible representations in $SL(2, \mathbb{C})$. arXiv:1605.08530, 2016.

E-mail address: s.sivek@imperial.ac.uk

IMPERIAL COLLEGE LONDON

E-mail address: raphael.zentner@mathematik.uni-regensburg.de

UNIVERSITÄT REGENSBURG